

PATENT COOPERATION TREATY

PCT

NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

Commissioner
 US Department of Commerce
 United States Patent and Trademark
 Office, PCT
 2011 South Clark Place Room
 CP2/5C24
 Arlington, VA 22202
 ETATS-UNIS D'AMERIQUE
 in its capacity as elected Office

Date of mailing (day/month/year) 07 February 2001 (07.02.01)	
International application No. PCT/EP00/05849	Applicant's or agent's file reference FB/BM45397
International filing date (day/month/year) 23 June 2000 (23.06.00)	Priority date (day/month/year) 25 June 1999 (25.06.99)
Applicant THONNARD, Joelle	

1. The designated Office is hereby notified of its election made:

☒ in the demand filed with the International Preliminary Examining Authority on:
 15 December 2000 (15.12.00)

☐ in a notice effecting later election filed with the International Bureau on:

2. The election ☒ was
☐ was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland Facsimile No.: (41-22) 740.14.35	Authorized officer A. Karkachi Telephone No.: (41-22) 338.83.38
------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

14


Applicant's or agent's file reference SD/FB/BM45397	FOR FURTHER ACTION See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. PCT/EP00/05849	International filing date (day/month/year) 23/06/2000	Priority date (day/month/year) 25/06/1999
International Patent Classification (IPC) or national classification and IPC C12N15/31		
Applicant SMITHKLINE BEECHAM BIOLOGICALS S.A. et al.		

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.
2. This REPORT consists of a total of 5 sheets, including this cover sheet.
 - ☐ This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☐ Priority
- III ☐ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
- IV ☐ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☒ Certain defects in the international application
- VIII ☐ Certain observations on the international application

Date of submission of the demand 15/12/2000	Date of completion of this report 28.08.2001
Name and mailing address of the international preliminary examining authority:  European Patent Office D-80298 Munich Tel. +49 89 2399 - 0 Tx: 523656 epmu d Fax: +49 89 2399 - 4465	Authorized officer Page, M Telephone No. +49 89 2399 7322



**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT**

International application No. PCT/EP00/05849

I. Basis of the report

1. With regard to the **elements** of the international application (*Replacement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to this report since they do not contain amendments (Rules 70.16 and 70.17)*):

Description, pages:

1-67 as originally filed

Claims, No.:

1-26 as originally filed

Drawings, sheets:

1/4-4/4 as originally filed

Sequence listing part of the description, pages:

1-2 (SEQ ID NOs. 1-6), as originally filed

2. With regard to the **language**, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language: , which is:

- ☐ the language of a translation furnished for the purposes of the international search (under Rule 23.1(b)).
- ☐ the language of publication of the international application (under Rule 48.3(b)).
- ☐ the language of a translation furnished for the purposes of international preliminary examination (under Rule 55.2 and/or 55.3).

3. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:

- ☒ contained in the international application in written form.
- ☒ filed together with the international application in computer readable form.
- ☐ furnished subsequently to this Authority in written form.
- ☐ furnished subsequently to this Authority in computer readable form.
- ☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
- ☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

4. The amendments have resulted in the cancellation of:

INTERNATIONAL PRELIMINARY
EXAMINATION REPORT

International application No. PCT/EP00/05849

- ☐ the description, pages:
☐ the claims, Nos.:
☐ the drawings, sheets:

5. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

(Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.)

6. Additional observations, if necessary:

V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

1. Statement

Novelty (N)	Yes:	Claims	1-26
	No:	Claims	
Inventive step (IS)	Yes:	Claims	
	No:	Claims	1-26
Industrial applicability (IA)	Yes:	Claims	1-26
	No:	Claims	

- 2. Citations and explanations**
see separate sheet

VII. Certain defects in the international application

The following defects in the form or contents of the international application have been noted:
see separate sheet

**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT - SEPARATE SHEET**

International application No. PCT/EP00/05849

The application concerns the provision of a novel outer membrane lipoprotein from *Moraxella catarrhalis* that is potentially of use in providing immunological protection against this organism.

Re Item V

Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

1) Reference is made to the following documents:

- D1: OCHSNER U A ET AL.: 'Pseudomonas aeruginosa fur overlaps with a gene encoding a novel outer membrane lipoprotein, OmlA' JOURNAL OF BACTERIOLOGY, vol. 181, no. 4, February 1999 (1999-02), pages 1099-1109, XP002154428
- D2: WO 94 10316 A (UNIV SASKATCHEWAN) 11 May 1994 (1994-05-11)
- D3: MURPHY T F: 'Branhamella catarrhalis: epidemiology, surface antigenic structure, and immune response' MICROBIOLOGICAL REVIEWS, vol. 60, no. 2, July 1996 (1996-07), pages 267-279, XP000857203 cited in the application
- D4: PANDHER K ET AL.: 'Genetic and immunologic analyses of PlpE, a lipoprotein important in complement-mediated killing of Pasteurella haemolytica serotype 1' INFECTION AND IMMUNITY, vol. 66, no. 12, December 1998 (1998-12), pages 5613-5619, XP002154430
- D5: GRAM T ET AL.: 'Improved diagnostic PCR assay for Actinobacillus pleuropneumoniae based on the nucleotide sequence of an outer membrane lipoprotein' JOURNAL OF CLINICAL MICROBIOLOGY, vol. 36, no. 2, February 1998 (1998-02), pages 443-448, XP002154429

2) **Novelty - Art.33(1) and (2) PCT:**

Claims 1-26 appear to be novel in light of the cited prior art. The closest polypeptide in the art is the OmlA protein of *Pseudomonas aeruginosa* disclosed in Fig. 1B of D1. This polypeptide is 31% identical to SEQ ID NOs. 2 and 4 of the present application.

3) **Inventive Step - Art.33(1) and (3) PCT:**

No specific prior art exists corresponding to the sequences provided by the application. However, in the absence of any concrete information regarding the function of the polypeptide of SEQ ID NOs. 2 or 4, the technical problem can only be regarded as the provision of an expressed polynucleotide sequence. Examples 5-11 concerning the possible functions of BASB112 appear to be purely theoretical in nature, rather than to demonstrate a proven function or use of the disclosed polypeptide. In order to recognise inventive step, the IPEA is obliged to satisfy itself that the problem underlying an application is indeed solved by the claimed compounds/methods.

The application fails to show whether BASB112 is a suitable vaccination protein as it has not been demonstrated that the protein is expressed, it has not been demonstrated that the protein is available as an antigen to the host organism (is it a cell-surface protein or not?) and it has not been demonstrated that the presence of antibodies against the protein are advantageous in protecting the host organism against infection by *M. catarrhalis*.

In light of this, the subject matter of **claims 1-26** cannot be considered to be inventive as no technical problem appears to be solved.

Re Item VII

Certain defects in the international application

- a) Contrary to the requirements of Rule 5.1(a)(ii) PCT, the relevant background art disclosed in the documents D1, D2, D4 and D5 is not mentioned in the description, nor are these document/s identified therein.

PATENT COOPERATION TREATY

From the
INTERNATIONAL PRELIMINARY EXAMINING AUTHORITY

To:

Peter John Giddings
SmithKline Beecham
Corporate Intellectual Property
Two New Horizons Court
Middlesex TW8 9EP
GRANDE BRETAGNE

PCT

NOTIFICATION OF TRANSMITTAL OF THE INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Rule 71.1)

Date of mailing
(day/month/year) 28.08.2001

Applicant's or agent's file reference
SD/FB/BM45397

IMPORTANT NOTIFICATION

International application No.
PCT/EP00/05849

International filing date (day/month/year)
23/06/2000

Priority date (day/month/year)
25/06/1999

Applicant
SMITHKLINE BEECHAM BIOLOGICALS S.A. et al.

1. The applicant is hereby notified that this International Preliminary Examining Authority transmits herewith the international preliminary examination report and its annexes, if any, established on the international application.
2. A copy of the report and its annexes, if any, is being transmitted to the International Bureau for communication to all the elected Offices.
3. Where required by any of the elected Offices, the International Bureau will prepare an English translation of the report (but not of any annexes) and will transmit such translation to those Offices.

4. REMINDER

The applicant must enter the national phase before each elected Office by performing certain acts (filing translations and paying national fees) within 30 months from the priority date (or later in some Offices) (Article 39(1)) (see also the reminder sent by the International Bureau with Form PCT/IB/301).

Where a translation of the international application must be furnished to an elected Office, that translation must contain a translation of any annexes to the international preliminary examination report. It is the applicant's responsibility to prepare and furnish such translation directly to each elected Office concerned.

For further details on the applicable time limits and requirements of the elected Offices, see Volume II of the PCT Applicant's Guide.

Name and mailing address of the IPEA/



European Patent Office
D-80298 Munich
Tel. +49 89 2399 - 0 Tx: 523656 epmu d
Fax: +49 89 2399 - 4465

Authorized officer


Büchler, S

Tel. +49 89 2399-8090



INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

Applicant's or agent's file reference SD/FB/BM45397		FOR FURTHER ACTION See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. PCT/EP00/05849	International filing date (day/month/year) 23/06/2000	Priority date (day/month/year) 25/06/1999	
International Patent Classification (IPC) or national classification and IPC C12N15/31			
Applicant SMITHKLINE BEECHAM BIOLOGICALS S.A. et al.			
<p>1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.</p> <p>2. This REPORT consists of a total of 5 sheets, including this cover sheet.</p> <p><input type="checkbox"/> This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).</p> <p>These annexes consist of a total of sheets.</p>			
<p>3. This report contains indications relating to the following items:</p> <p>I <input checked="" type="checkbox"/> Basis of the report</p> <p>II <input type="checkbox"/> Priority</p> <p>III <input type="checkbox"/> Non-establishment of opinion with regard to novelty, inventive step and industrial applicability</p> <p>IV <input type="checkbox"/> Lack of unity of invention</p> <p>V <input checked="" type="checkbox"/> Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement</p> <p>VI <input type="checkbox"/> Certain documents cited</p> <p>VII <input checked="" type="checkbox"/> Certain defects in the international application</p> <p>VIII <input type="checkbox"/> Certain observations on the international application</p>			
Date of submission of the demand 15/12/2000		Date of completion of this report 28.08.2001	
Name and mailing address of the international preliminary examining authority:  European Patent Office D-80298 Munich Tel. +49 89 2399 - 0 Tx: 523656 epmu d Fax: +49 89 2399 - 4465		Authorized officer Page, M Telephone No. +49 89 2399 7322	



**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT**

International application No. PCT/EP00/05849

I. Basis of the report

1. With regard to the **elements** of the international application (*Replacement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to this report since they do not contain amendments (Rules 70.16 and 70.17)*):

Description, pages:

1-67 as originally filed

Claims, No.:

1-26 as originally filed

Drawings, sheets:

1/4-4/4 as originally filed

Sequence listing part of the description, pages:

1-2 (SEQ ID NOs. 1-6), as originally filed

2. With regard to the **language**, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language: , which is:

- ☐ the language of a translation furnished for the purposes of the international search (under Rule 23.1(b)).
- ☐ the language of publication of the international application (under Rule 48.3(b)).
- ☐ the language of a translation furnished for the purposes of international preliminary examination (under Rule 55.2 and/or 55.3).

3. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:

- ☒ contained in the international application in written form.
- ☒ filed together with the international application in computer readable form.
- ☐ furnished subsequently to this Authority in written form.
- ☐ furnished subsequently to this Authority in computer readable form.
- ☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
- ☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

4. The amendments have resulted in the cancellation of:

**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT**

International application No. PCT/EP00/05849

- ☐ the description, pages:
☐ the claims, Nos.:
☐ the drawings, sheets:

5. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

(Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.)

6. Additional observations, if necessary:

V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

1. Statement

Novelty (N)	Yes:	Claims	1-26
	No:	Claims	
Inventive step (IS)	Yes:	Claims	
	No:	Claims	1-26
Industrial applicability (IA)	Yes:	Claims	1-26
	No:	Claims	

- 2. Citations and explanations**
see separate sheet

VII. Certain defects in the international application

The following defects in the form or contents of the international application have been noted:
see separate sheet

The application concerns the provision of a novel outer membrane lipoprotein from *Moraxella catarrhalis* that is potentially of use in providing immunological protection against this organism.

Re Item V

Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

1) Reference is made to the following documents:

- D1: OCHSNER U A ET AL.: 'Pseudomonas aeruginosa fur overlaps with a gene encoding a novel outer membrane lipoprotein, OmlA' JOURNAL OF BACTERIOLOGY, vol. 181, no. 4, February 1999 (1999-02), pages 1099-1109, XP002154428
- D2: WO 94 10316 A (UNIV SASKATCHEWAN) 11 May 1994 (1994-05-11)
- D3: MURPHY T F: 'Branhamella catarrhalis: epidemiology, surface antigenic structure, and immune response' MICROBIOLOGICAL REVIEWS, vol. 60, no. 2, July 1996 (1996-07), pages 267-279, XP000857203 cited in the application
- D4: PANDHER K ET AL.: 'Genetic and immunologic analyses of PlpE, a lipoprotein important in complement-mediated killing of Pasteurella haemolytica serotype 1' INFECTION AND IMMUNITY, vol. 66, no. 12, December 1998 (1998-12), pages 5613-5619, XP002154430
- D5: GRAM T ET AL.: 'Improved diagnostic PCR assay for Actinobacillus pleuropneumoniae based on the nucleotide sequence of an outer membrane lipoprotein' JOURNAL OF CLINICAL MICROBIOLOGY, vol. 36, no. 2, February 1998 (1998-02), pages 443-448, XP002154429

2) **Novelty - Art.33(1) and (2) PCT:**

Claims 1-26 appear to be novel in light of the cited prior art. The closest polypeptide in the art is the OmlA protein of *Pseudomonas aeruginosa* disclosed in Fig. 1B of D1. This polypeptide is 31% identical to SEQ ID NOs. 2 and 4 of the present application.

3) **Inventive Step - Art.33(1) and (3) PCT:**

No specific prior art exists corresponding to the sequences provided by the application. However, in the absence of any concrete information regarding the function of the polypeptide of SEQ ID NOs. 2 or 4, the technical problem can only be regarded as the provision of an expressed polynucleotide sequence. Examples 5-11 concerning the possible functions of BASB112 appear to be purely theoretical in nature, rather than to demonstrate a proven function or use of the disclosed polypeptide. In order to recognise inventive step, the IPEA is obliged to satisfy itself that the problem underlying an application is indeed solved by the claimed compounds/methods.

The application fails to show whether BASB112 is a suitable vaccination protein as it has not been demonstrated that the protein is expressed, it has not been demonstrated that the protein is available as an antigen to the host organism (is it a cell-surface protein or not?) and it has not been demonstrated that the presence of antibodies against the protein are advantageous in protecting the host organism against infection by *M. catarrhalis*.

In light of this, the subject matter of **claims 1-26** cannot be considered to be inventive as no technical problem appears to be solved.

Re Item VII

Certain defects in the international application

- a) Contrary to the requirements of Rule 5.1(a)(ii) PCT, the relevant background art disclosed in the documents D1, D2, D4 and D5 is not mentioned in the description, nor are these document/s identified therein.

PATENT COOPERATION TREATY

From the INTERNATIONAL SEARCHING AUTHORITY

PCT

NOTIFICATION OF TRANSMITTAL OF THE INTERNATIONAL SEARCH REPORT OR THE DECLARATION

(PCT Rule 44.1)

To:

SmithKline Beecham
Corporate Intellectual Property
Attn. PRIVETT, Kathryn L.
Two New Horizons Court
Brentford
Middlesex TW8 9EP
UNITED KINGDOM

Date of mailing
(day/month/year)

15/12/2000

Applicant's or agent's file reference

FB/BM45397

FOR FURTHER ACTION

See paragraphs 1 and 4 below

International application No.

PCT/EP 00/05849

International filing date
(day/month/year)

23/06/2000

Applicant

SMITHKLINE BEECHAM BIOLOGICALS S.A.

1. ☒ The applicant is hereby notified that the International Search Report has been established and is transmitted herewith.

Filing of amendments and statement under Article 19:

The applicant is entitled, if he so wishes, to amend the claims of the International Application (see Rule 46):

When? The time limit for filing such amendments is normally 2 months from the date of transmittal of the International Search Report; however, for more details, see the notes on the accompanying sheet.

Where? Directly to the International Bureau of WIPO
34, chemin des Colombettes
1211 Geneva 20, Switzerland
Facsimile No.: (41-22) 740.14.35

For more detailed instructions, see the notes on the accompanying sheet.

2. ☐ The applicant is hereby notified that no International Search Report will be established and that the declaration under Article 17(2)(a) to that effect is transmitted herewith.

3. ☐ **With regard to the protest** against payment of (an) additional fee(s) under Rule 40.2, the applicant is notified that:

☐ the protest together with the decision thereon has been transmitted to the International Bureau together with the applicant's request to forward the texts of both the protest and the decision thereon to the designated Offices.

☐ no decision has been made yet on the protest; the applicant will be notified as soon as a decision is made.

4. **Further action(s):** The applicant is reminded of the following:

Shortly after **18 months** from the priority date, the international application will be published by the International Bureau. If the applicant wishes to avoid or postpone publication, a notice of withdrawal of the international application, or of the priority claim, must reach the International Bureau as provided in Rules 90bis.1 and 90bis.3, respectively, before the completion of the technical preparations for international publication.

Within **19 months** from the priority date, a demand for international preliminary examination must be filed if the applicant wishes to postpone the entry into the national phase until 30 months from the priority date (in some Offices even later).

Within **20 months** from the priority date, the applicant must perform the prescribed acts for entry into the national phase before all designated Offices which have not been elected in the demand or in a later election within 19 months from the priority date or could not be elected because they are not bound by Chapter II.

Name and mailing address of the International Searching Authority



European Patent Office, P.B. 5818 Patentlaan 2
NL-2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Mireille Claudepierre

NOTES TO FORM PCT/ISA/220

These Notes are intended to give the basic instructions concerning the filing of amendments under article 19. The Notes are based on the requirements of the Patent Cooperation Treaty, the Regulations and the Administrative Instructions under that Treaty. In case of discrepancy between these Notes and those requirements, the latter are applicable. For more detailed information, see also the PCT Applicant's Guide, a publication of WIPO.

In these Notes, "Article", "Rule", and "Section" refer to the provisions of the PCT, the PCT Regulations and the PCT Administrative Instructions respectively.

INSTRUCTIONS CONCERNING AMENDMENTS UNDER ARTICLE 19

The applicant has, after having received the international search report, one opportunity to amend the claims of the international application. It should however be emphasized that, since all parts of the international application (claims, description and drawings) may be amended during the international preliminary examination procedure, there is usually no need to file amendments of the claims under Article 19 except where, e.g. the applicant wants the latter to be published for the purposes of provisional protection or has another reason for amending the claims before international publication. Furthermore, it should be emphasized that provisional protection is available in some States only.

What parts of the international application may be amended?

Under Article 19, only the claims may be amended.

During the international phase, the claims may also be amended (or further amended) under Article 34 before the International Preliminary Examining Authority. The description and drawings may only be amended under Article 34 before the International Examining Authority.

Upon entry into the national phase, all parts of the international application may be amended under Article 28 or, where applicable, Article 41.

When?

Within 2 months from the date of transmittal of the international search report or 16 months from the priority date, whichever time limit expires later. It should be noted, however, that the amendments will be considered as having been received on time if they are received by the International Bureau after the expiration of the applicable time limit but before the completion of the technical preparations for international publication (Rule 46.1).

Where not to file the amendments?

The amendments may only be filed with the International Bureau and not with the receiving Office or the International Searching Authority (Rule 46.2).

Where a demand for international preliminary examination has been/is filed, see below.

How?

Either by cancelling one or more entire claims, by adding one or more new claims or by amending the text of one or more of the claims as filed.

A replacement sheet must be submitted for each sheet of the claims which, on account of an amendment or amendments, differs from the sheet originally filed.

All the claims appearing on a replacement sheet must be numbered in Arabic numerals. Where a claim is cancelled, no renumbering of the other claims is required. In all cases where claims are renumbered, they must be renumbered consecutively (Administrative Instructions, Section 205(b)).

The amendments must be made in the language in which the international application is to be published.

What documents must/may accompany the amendments?

Letter (Section 205(b)):

The amendments must be submitted with a letter.

The letter will not be published with the international application and the amended claims. It should not be confused with the "Statement under Article 19(1)" (see below, under "Statement under Article 19(1)").

The letter must be in English or French, at the choice of the applicant. However, if the language of the international application is English, the letter must be in English; if the language of the international application is French, the letter must be in French.

NOTES TO FORM PCT/ISA/220 (c ntinued)

The letter must indicate the differences between the claims as filed and the claims as amended. It must, in particular, indicate, in connection with each claim appearing in the international application (it being understood that identical indications concerning several claims may be grouped), whether

- (i) the claim is unchanged;
- (ii) the claim is cancelled;
- (iii) the claim is new;
- (iv) the claim replaces one or more claims as filed;
- (v) the claim is the result of the division of a claim as filed.

The following examples illustrate the manner in which amendments must be explained in the accompanying letter:

1. [Where originally there were 48 claims and after amendment of some claims there are 51]:
"Claims 1 to 29, 31, 32, 34, 35, 37 to 48 replaced by amended claims bearing the same numbers; claims 30, 33 and 36 unchanged; new claims 49 to 51 added."
2. [Where originally there were 15 claims and after amendment of all claims there are 11]:
"Claims 1 to 15 replaced by amended claims 1 to 11."
3. [Where originally there were 14 claims and the amendments consist in cancelling some claims and in adding new claims]:
"Claims 1 to 6 and 14 unchanged; claims 7 to 13 cancelled; new claims 15, 16 and 17 added." or
"Claims 7 to 13 cancelled; new claims 15, 16 and 17 added; all other claims unchanged."
4. [Where various kinds of amendments are made]:
"Claims 1-10 unchanged; claims 11 to 13, 18 and 19 cancelled; claims 14, 15 and 16 replaced by amended claim 14; claim 17 subdivided into amended claims 15, 16 and 17; new claims 20 and 21 added."

"Statement under article 19(1)" (Rule 46.4)

The amendments may be accompanied by a statement explaining the amendments and indicating any impact that such amendments might have on the description and the drawings (which cannot be amended under Article 19(1)).

The statement will be published with the international application and the amended claims.

It must be in the language in which the international application is to be published.

It must be brief, not exceeding 500 words if in English or if translated into English.

It should not be confused with and does not replace the letter indicating the differences between the claims as filed and as amended. It must be filed on a separate sheet and must be identified as such by a heading, preferably by using the words "Statement under Article 19(1)."

It may not contain any disparaging comments on the international search report or the relevance of citations contained in that report. Reference to citations, relevant to a given claim, contained in the international search report may be made only in connection with an amendment of that claim.

Consequence if a demand for international preliminary examination has already been filed

If, at the time of filing any amendments under Article 19, a demand for international preliminary examination has already been submitted, the applicant must preferably, at the same time of filing the amendments with the International Bureau, also file a copy of such amendments with the International Preliminary Examining Authority (see Rule 62.2(a), first sentence).

Consequence with regard to translation of the international application for entry into the national phase

The applicant's attention is drawn to the fact that, where upon entry into the national phase, a translation of the claims as amended under Article 19 may have to be furnished to the designated/elected Offices, instead of, or in addition to, the translation of the claims as filed.

For further details on the requirements of each designated/elected Office, see Volume II of the PCT Applicant's Guide.

INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference FB/BM45397	FOR FURTHER ACTION see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.	
International application No. PCT/EP 00/ 05849	International filing date (day/month/year) 23/06/2000	(Earliest) Priority Date (day/month/year) 25/06/1999
Applicant SMITHKLINE BEECHAM BIOLOGICALS S.A.		

This International Search Report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This International Search Report consists of a total of 3 sheets.

☒ It is also accompanied by a copy of each prior art document cited in this report.

1. Basis of the report

- a. With regard to the **language**, the international search was carried out on the basis of the international application in the language in which it was filed, unless otherwise indicated under this item.

☐ the international search was carried out on the basis of a translation of the international application furnished to this Authority (Rule 23.1(b)).

- b. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international search was carried out on the basis of the sequence listing :

☒ contained in the international application in written form.

☒ filed together with the international application in computer readable form.

☐ furnished subsequently to this Authority in written form.

☐ furnished subsequently to this Authority in computer readable form.

☐ the statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.

☐ the statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished

2. ☐ **Certain claims were found unsearchable** (See Box I).

3. ☐ **Unity of invention is lacking** (see Box II).

4. With regard to the **title**,

☐ the text is approved as submitted by the applicant.

☒ the text has been established by this Authority to read as follows:

MORAXELLA CATARRHALIS POLYPEPTIDES

5. With regard to the **abstract**,

☒ the text is approved as submitted by the applicant.

☐ the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this international search report, submit comments to this Authority.

6. The figure of the **drawings** to be published with the abstract is Figure No.

☐ as suggested by the applicant.

☐ because the applicant failed to suggest a figure.

☐ because this figure better characterizes the invention.

☒ None of the figures.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/05849

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/31 C12N15/62 C07K14/21 C07K16/12 A61K39/02
 A61K48/00 G01N33/50

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K A61K G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EMBL, STRAND, WPI Data, EPO-Internal, PAJ, BIOSIS, MEDLINE, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	OCHSNER U A ET AL.: "Pseudomonas aeruginosa fur overlaps with a gene encoding a novel outer membrane lipoprotein, OmlA" JOURNAL OF BACTERIOLOGY, vol. 181, no. 4, February 1999 (1999-02), pages 1099-1109, XP002154428 Note: 31 % aa seq identity with SEQ ID NO:2 in 116 aa overlap. abstract	1-5,7-13
A	WO 94 10316 A (UNIV SASKATCHEWAN) 11 May 1994 (1994-05-11) claims 1-11; examples 1-6 --- -/--	1-26



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

1 December 2000

Date of mailing of the international search report

15/12/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

van de Kamp, M

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 00/05849

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	MURPHY T F: "Branhamella catarrhalis: epidemiology, surface antigenic structure, and immune response" MICROBIOLOGICAL REVIEWS, vol. 60, no. 2, July 1996 (1996-07), pages 267-279, XP000857203 cited in the application page 271, right-hand column, line 57 -page 273, right-hand column, line 27; figure 2; table 3 ---	1-26
A	WO 98 55606 A (LOOSMORE SHEENA M ;CONNAUGHT LAB (CA); WANG QUIJUN (CA); YANG YAN) 10 December 1998 (1998-12-10) page 3, line 20 -page 11, line 5 example 4 ---	1-26
A	PANDHER K ET AL.: "Genetic and immunologic analyses of PlpE, a lipoprotein important in complement-mediated killing of Pasteurella haemolytica serotype 1" INFECTION AND IMMUNITY, vol. 66, no. 12, December 1998 (1998-12), pages 5613-5619, XP002154430 abstract ---	19-22
A	GRAM T ET AL.: "Improved diagnostic PCR assay for Actinobacillus pleuropneumoniae based on the nucleotide sequence of an outer membrane lipoprotein" JOURNAL OF CLINICAL MICROBIOLOGY, vol. 36, no. 2, February 1998 (1998-02), pages 443-448, XP002154429 abstract -----	23

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 00/05849

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9410316 A	11-05-1994	US 5441736 A CA 2148369 A EP 0669984 A US 5891677 A	15-08-1995 11-05-1994 06-09-1995 06-04-1999
WO 9855606 A	10-12-1998	US 5977337 A AU 7753498 A EP 1000144 A	02-11-1999 21-12-1998 17-05-2000

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
4 January 2001 (04.01.2001)

PCT

(10) International Publication Number
WO 01/00835 A1

(51) International Patent Classification⁷: C12N 15/31,
15/62, C07K 14/21, 16/12, A61K 39/02, 48/00, G01N
33/50

(21) International Application Number: PCT/EP00/05849

(22) International Filing Date: 23 June 2000 (23.06.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
9914870.2 25 June 1999 (25.06.1999) GB

(71) Applicant (for all designated States except US):
SMITHKLINE BEECHAM BIOLOGICALS S.A.
[BE/BE]; Rue de l'Institut 89, B-1330 Rixensart (BE).

(72) Inventor; and

(75) Inventor/Applicant (for US only): THONNARD, Joelle
[BE/BE]; SmithKline Beecham Biologicals s.a., Rue de
L'Institut 89, B-1330 Rixensart (BE).

(74) Agent: PRIVETT, Kathryn, Louise; SmithKline
Beecham, Corporate Intellectual Property, Two New
Horizons Court, Brentford, Middlesex TW8 9EP (GB).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- With international search report.
- Before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments.

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: MORAXELLA CATARRHALIS POLYPEPTIDES

(57) Abstract: The invention provides BASB112 polypeptides and polynucleotides encoding BASB112 polypeptides and methods for producing such polypeptides by recombinant techniques. Also provided are diagnostic, prophylactic and therapeutic uses.

13 DEC 2001

WO 01/00835 A1

THE
WORLD
OF
THE
FUTURE

MORAXELLA CATARRHALIS POLYPEPTIDES

FIELD OF THE INVENTION

This invention relates to polynucleotides, (herein referred to as "BASB112 polynucleotide(s)"), polypeptides encoded by them (referred to herein as "BASB112" or "BASB112 polypeptide(s)"), recombinant materials and methods for their production. In another aspect, the invention relates to methods for using such polypeptides and polynucleotides, including vaccines against bacterial infections. In a further aspect, the invention relates to diagnostic assays for detecting infection of certain pathogens.

BACKGROUND OF THE INVENTION

Moraxella catarrhalis (also named *Branhamella catarrhalis*) is a Gram negative bacteria frequently isolated from the human upper respiratory tract. It is responsible for several pathologies the main ones being otitis media in infants and children, and pneumonia in elderlies. It is also responsible of sinusitis, nosocomial infections and less frequently of invasive diseases.

Otitis media is an important childhood disease both by the number of cases and its potential sequelae. More than 3.5 millions cases are recorded every year in the United States, and it is estimated that 80 % of the children have experienced at least one episode of otitis before reaching the age of 3 (Klein, JO (1994) Clin.Inf.Dis 19:823). Left untreated, or becoming chronic, this disease may lead to hearing losses that could be temporary (in the case of fluid accumulation in the middle ear) or permanent (if the auditive nerve is damaged). In infants, such hearing losses may be responsible for a delayed speech learning.

Three bacterial species are primarily isolated from the middle ear of children with otitis media: *Streptococcus pneumoniae*, non typeable *Haemophilus influenza* (NTHi) and *M. catarrhalis*. They are present in 60 to 90 % of the cases. A review of recent studies shows that *S. pneumoniae* and NTHi represent both about 30 %, and *M. catarrhalis* about 15 % of

the otitis media cases (Murphy, TF (1996) Microbiol.Rev. 60:267). Other bacteria could be isolated from the middle ear (*H. influenza* type B, *S. pyogenes* etc) but at a much lower frequency (2 % of the cases or less).

Epidemiological data indicate that, for the pathogens found in the middle ear, the colonization of the upper respiratory tract is an absolute prerequisite for the development of an otitis; other are however also required to lead to the disease (Dickinson, DP et al. (1988) J. Infect.Dis. 158:205, Faden, HL et al. (1991) Ann.Otorhinol.Laryngol. 100:612). These are important to trigger the migration of the bacteria into the middle ear via the Eustachian tubes, followed by the initiation of an inflammatory process. These factors are unknown todate. It has been postulated that a transient anomaly of the immune system following a viral infection, for example, could cause an inability to control the colonization of the respiratory tract (Faden, HL et al (1994) J. Infect.Dis. 169:1312). An alternative explanation is that the exposure to environmental factors allow a more important colonization of some children, who subsequently become susceptible to the development of otitis media because of the sustained presence of middle ear pathogens (Murphy, TF (1996) Microbiol.Rev. 60:267).

The immune response to *M. catarrhalis* is poorly characterized. The analysis of strains isolated sequentially from the nasopharynx of babies followed from 0 to 2 years of age, indicates that they get and eliminate frequently new strains. This indicates that an efficacious immune response against this bacteria is mounted by the colonized children (Faden, HL et al (1994) J. Infect.Dis. 169:1312).

In most adults tested, bactericidal antibodies have been identified (Chapman, AJ et al. (1985) J. Infect.Dis. 151:878). Strains of *M. catarrhalis* present variations in their capacity to resist serum bactericidal activity: in general, isolates from diseased individuals are more resistant than those who are simply colonized (Hol, C et al. (1993) Lancet 341:1281, Jordan, KL et al. (1990) Am.J.Med. 88 (suppl. 5A):28S). Serum resistance could therefore be

considered as a virulence factor of the bacteria. An opsonizing activity has been observed in the sera of children recovering from otitis media.

The antigens targetted by these different immune responses in humans have not been identified, with the exception of OMP B1, a 84 kDa protein which expression is regulated by iron, and that is recognized by the sera of patients with pneumonia (Sethi, S, et al. (1995) Infect.Immun. 63:1516) , and of UspA1 and UspA2 (Chen D. et al.(1999), Infect.Immun. 67:1310).

A few other membrane proteins present on the surface of *M. catarrhalis* have been characterized using biochemical method, or for their potential implication in the induction of a protective immunity (for review, see Murphy, TF (1996) Microbiol.Rev. 60:267). In a mouse pneumonia model, the presence of antibodies raised against some of them (UspA, CopB) favors a faster clearance of the pulmonary infection. Another polypeptide (OMP CD) is highly conserved among *M. catarrhalis* strains, and presents homologies with a porin of *Pseudomonas aeruginosa*, which has been demonstrated efficacious against this bacterium in animal models.

The frequency of *Moraxella catarrhalis* infections has risen dramatically in the past few decades. This has been attributed to the emergence of multiply antibiotic resistant strains and an increasing population of people with weakened immune systems. It is no longer uncommon to isolate *Moraxella catarrhalis* strains that are resistant to some or all of the standard antibiotics. This phenomenon has created an unmet medical need and demand for new anti-microbial agents, vaccines, drug screening methods, and diagnostic tests for this organism.

SUMMARY OF THE INVENTION

The present invention relates to BASB112, in particular BASB112 polypeptides and BASB112 polynucleotides, recombinant materials and methods for their production. In another aspect, the invention relates to methods for using such polypeptides and polynucleotides, including prevention and treatment of microbial diseases, amongst others. In a further aspect, the invention relates to diagnostic assays for detecting diseases associated with microbial infections and conditions associated with such infections, such as assays for detecting expression or activity of BASB112 polynucleotides or polypeptides.

Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following descriptions and from reading the other parts of the present disclosure.

DESCRIPTION OF THE INVENTION

The invention relates to BASB112 polypeptides and polynucleotides as described in greater detail below. In particular, the invention relates to polypeptides and polynucleotides of BASB112 of *Moraxella catarrhalis*, which have the characteristics of a lipoprotein (or a polynucleotide encoding a lipoprotein). BASB112 polypeptides are thus surface exposed proteins and useful vaccine candidates. BASB112 is found to be related by amino acid sequence homology to *Pseudomonas aeruginosa* lipoprotein (OmlA). The invention relates especially to BASB112 having the nucleotide and amino acid sequences set out in SEQ ID NO:1 or 3 and SEQ ID NO:2 or 4 respectively. It is understood that sequences recited in the Sequence Listing below as "DNA" represent an exemplification of one embodiment of the invention, since those of ordinary skill will recognize that such sequences can be usefully employed in polynucleotides in general, including ribopolynucleotides.

Polypeptides

In one aspect of the invention there are provided polypeptides of *Moraxella catarrhalis* referred to herein as "BASB112" and "BASB112 polypeptides" as well as biologically,

diagnostically, prophylactically, clinically or therapeutically useful variants thereof, and compositions comprising the same.

The present invention further provides for:

- (a) an isolated polypeptide which comprises an amino acid sequence which has at least 85% identity, preferably at least 90% identity, more preferably at least 95% identity, most preferably at least 97-99% or exact identity, to that of SEQ ID NO:2 or 4;
- (b) a polypeptide encoded by an isolated polynucleotide comprising a polynucleotide sequence which has at least 85% identity, preferably at least 90% identity, more preferably at least 95% identity, even more preferably at least 97-99% or exact identity to SEQ ID NO:1 or 3 over the entire length of SEQ ID NO:1 or 3 respectively; or
- (c) a polypeptide encoded by an isolated polynucleotide comprising a polynucleotide sequence encoding a polypeptide which has at least 85% identity, preferably at least 90% identity, more preferably at least 95% identity, even more preferably at least 97-99% or exact identity, to the amino acid sequence of SEQ ID NO:2 or 4.

The BASB112 polypeptides provided in SEQ ID NO:2 or 4 are the BASB112 polypeptides from *Moraxella catarrhalis* strain Mc2931 (ATCC 43617).

The invention also provides an immunogenic fragment of a BASB112 polypeptide, that is, a contiguous portion of the BASB112 polypeptide which has the same or substantially the same immunogenic activity as the polypeptide comprising the amino acid sequence of SEQ ID NO:2 or 4; That is to say, the fragment (if necessary when coupled to a carrier) is capable of raising an immune response which recognises the BASB112 polypeptide. Such an immunogenic fragment may include, for example, the BASB112 polypeptide lacking an N-terminal leader sequence, and/or a transmembrane domain and/or a C-terminal anchor domain. In a preferred aspect the immunogenic fragment of BASB112 according to the invention comprises substantially all of the extracellular domain of a polypeptide which has at least 85% identity, preferably at least 90% identity, more

preferably at least 95% identity, most preferably at least 97-99% identity, to that of SEQ ID NO:2 or 4 over the entire length of SEQ ID NO:2 or 4 respectively.

A fragment is a polypeptide having an amino acid sequence that is entirely the same as part but not all of any amino acid sequence of any polypeptide of the invention. As with BASB112 polypeptides, fragments may be "free-standing," or comprised within a larger polypeptide of which they form a part or region, most preferably as a single continuous region in a single larger polypeptide.

Preferred fragments include, for example, truncation polypeptides having a portion of an amino acid sequence of SEQ ID NO:2 or 4 or of variants thereof, such as a continuous series of residues that includes an amino- and/or carboxyl-terminal amino acid sequence. Degradation forms of the polypeptides of the invention produced by or in a host cell, are also preferred. Further preferred are fragments characterized by structural or functional attributes such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions.

Further preferred fragments include an isolated polypeptide comprising an amino acid sequence having at least 15, 20, 30, 40, 50 or 100 contiguous amino acids from the amino acid sequence of SEQ ID NO:2 or 4, or an isolated polypeptide comprising an amino acid sequence having at least 15, 20, 30, 40, 50 or 100 contiguous amino acids truncated or deleted from the amino acid sequence of SEQ ID NO:2 or 4.

Fragments of the polypeptides of the invention may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, these fragments

may be employed as intermediates for producing the full-length polypeptides of the invention.

Particularly preferred are variants in which several, 5-10, 1-5, 1-3, 1-2 or 1 amino acids are substituted, deleted, or added in any combination.

The polypeptides, or immunogenic fragments, of the invention may be in the form of the "mature" protein or may be a part of a larger protein such as a precursor or a fusion protein. It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, pro-sequences, sequences which aid in purification such as multiple histidine residues, or an additional sequence for stability during recombinant production. Furthermore, addition of exogenous polypeptide or lipid tail or polynucleotide sequences to increase the immunogenic potential of the final molecule is also considered.

In one aspect, the invention relates to genetically engineered soluble fusion proteins comprising a polypeptide of the present invention, or a fragment thereof, and various portions of the constant regions of heavy or light chains of immunoglobulins of various subclasses (IgG, IgM, IgA, IgE). Preferred as an immunoglobulin is the constant part of the heavy chain of human IgG, particularly IgG1, where fusion takes place at the hinge region. In a particular embodiment, the Fc part can be removed simply by incorporation of a cleavage sequence which can be cleaved with blood clotting factor Xa.

Furthermore, this invention relates to processes for the preparation of these fusion proteins by genetic engineering, and to the use thereof for drug screening, diagnosis and therapy. A further aspect of the invention also relates to polynucleotides encoding such fusion proteins. Examples of fusion protein technology can be found in International Patent Application Nos. WO94/29458 and WO94/22914.

The proteins may be chemically conjugated, or expressed as recombinant fusion proteins allowing increased levels to be produced in an expression system as compared to non-fused protein. The fusion partner may assist in providing T helper epitopes (immunological fusion partner), preferably T helper epitopes recognised by humans, or assist in expressing the protein (expression enhancer) at higher yields than the native recombinant protein. Preferably the fusion partner will be both an immunological fusion partner and expression enhancing partner.

Fusion partners include protein D from *Haemophilus influenzae* and the non-structural protein from influenzae virus, NS1 (hemagglutinin). Another fusion partner is the protein known as LytA. Preferably the C terminal portion of the molecule is used. LytA is derived from *Streptococcus pneumoniae* which synthesize an N-acetyl-L-alanine amidase, amidase LytA, (coded by the *lytA* gene {Gene, 43 (1986) page 265-272}) an autolysin that specifically degrades certain bonds in the peptidoglycan backbone. The C-terminal domain of the LytA protein is responsible for the affinity to the choline or to some choline analogues such as DEAE. This property has been exploited for the development of E.coli C-LytA expressing plasmids useful for expression of fusion proteins. Purification of hybrid proteins containing the C-LytA fragment at its amino terminus has been described {Biotechnology: 10, (1992) page 795-798}. It is possible to use the repeat portion of the LytA molecule found in the C terminal end starting at residue 178, for example residues 188 - 305.

The present invention also includes variants of the aforementioned polypeptides, that is polypeptides that vary from the referents by conservative amino acid substitutions, whereby a residue is substituted by another with like characteristics. Typical such substitutions are among Ala, Val, Leu and Ile; among Ser and Thr; among the acidic residues Asp and Glu; among Asn and Gln; and among the basic residues Lys and Arg; or aromatic residues Phe and Tyr.

Polypeptides of the present invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

It is most preferred that a polypeptide of the invention is derived from *Moraxella catarrhalis*, however, it may preferably be obtained from other organisms of the same taxonomic genus. A polypeptide of the invention may also be obtained, for example, from organisms of the same taxonomic family or order.

Polynucleotides

It is an object of the invention to provide polynucleotides that encode BASB112 polypeptides, particularly polynucleotides that encode the polypeptide herein designated BASB112.

In a particularly preferred embodiment of the invention the polynucleotide comprises a region encoding BASB112 polypeptides comprising a sequence set out in SEQ ID NO:1 or 3 which includes a full length gene, or a variant thereof.

The BASB112 polynucleotides provided in SEQ ID NO:1 or 3 are the BASB112 polynucleotides from *Moraxella catarrhalis* strain Mc2931 (ATCC 43617).

As a further aspect of the invention there are provided isolated nucleic acid molecules encoding and/or expressing BASB112 polypeptides and polynucleotides, particularly *Moraxella catarrhalis* BASB112 polypeptides and polynucleotides, including, for example, unprocessed RNAs, ribozyme RNAs, mRNAs, cDNAs, genomic DNAs, B- and Z-DNAs. Further embodiments of the invention include biologically,

diagnostically, prophylactically, clinically or therapeutically useful polynucleotides and polypeptides, and variants thereof, and compositions comprising the same.

Another aspect of the invention relates to isolated polynucleotides, including at least one full length gene, that encodes a BASB112 polypeptide having a deduced amino acid sequence of SEQ ID NO:2 or 4 and polynucleotides closely related thereto and variants thereof.

In another particularly preferred embodiment of the invention there is a BASB112 polypeptide from *Moraxella catarrhalis* comprising or consisting of an amino acid sequence of SEQ ID NO:2 or 4 or a variant thereof.

Using the information provided herein, such as a polynucleotide sequence set out in SEQ ID NO:1 or 3, a polynucleotide of the invention encoding BASB112 polypeptide may be obtained using standard cloning and screening methods, such as those for cloning and sequencing chromosomal DNA fragments from bacteria using *Moraxella catarrhalis* Catlin cells as starting material, followed by obtaining a full length clone. For example, to obtain a polynucleotide sequence of the invention, such as a polynucleotide sequence given in SEQ ID NO:1 or 3, typically a library of clones of chromosomal DNA of *Moraxella catarrhalis* Catlin in *E.coli* or some other suitable host is probed with a radiolabeled oligonucleotide, preferably a 17-mer or longer, derived from a partial sequence. Clones carrying DNA identical to that of the probe can then be distinguished using stringent hybridization conditions. By sequencing the individual clones thus identified by hybridization with sequencing primers designed from the original polypeptide or polynucleotide sequence it is then possible to extend the polynucleotide sequence in both directions to determine a full length gene sequence. Conveniently, such sequencing is performed, for example, using denatured double stranded DNA prepared from a plasmid clone. Suitable techniques are described by Maniatis, T., Fritsch, E.F. and Sambrook et al., *MOLECULAR CLONING, A LABORATORY MANUAL*, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989). (see in particular Screening By

Hybridization 1.90 and Sequencing Denatured Double-Stranded DNA Templates 13.70). Direct genomic DNA sequencing may also be performed to obtain a full length gene sequence. Illustrative of the invention, each polynucleotide set out in SEQ ID NO:1 or 3 was discovered in a DNA library derived from *Moraxella catarrhalis*.

Moreover, each DNA sequence set out in SEQ ID NO:1 or 3 contains an open reading frame encoding a protein having about the number of amino acid residues set forth in SEQ ID NO:2 or 4 with a deduced molecular weight that can be calculated using amino acid residue molecular weight values well known to those skilled in the art.

The polynucleotide of SEQ ID NO:1, between the start codon at nucleotide number 1 and the stop codon which begins at nucleotide number 367 of SEQ ID NO:1, encodes the polypeptide of SEQ ID NO:2.

The polynucleotide of SEQ ID NO:3, between the start codon at nucleotide number 1 and the stop codon which begins at nucleotide number 364 of SEQ ID NO:3, encodes the polypeptide of SEQ ID NO:4.

In a further aspect, the present invention provides for an isolated polynucleotide comprising or consisting of:

- (a) a polynucleotide sequence which has at least 85% identity, preferably at least 90% identity, more preferably at least 95% identity, even more preferably at least 97-99% or exact identity to SEQ ID NO:1 or 3 over the entire length of SEQ ID NO:1 or 3 respectively; or
- (b) a polynucleotide sequence encoding a polypeptide which has at least 85% identity, preferably at least 90% identity, more preferably at least 95% identity, even more preferably at least 97-99% or 100% exact, to the amino acid sequence of SEQ ID NO:2 or 4, over the entire length of SEQ ID NO:2 or 4 respectively.

A polynucleotide encoding a polypeptide of the present invention, including homologs and orthologs from species other than *Moraxella catarrhalis*, may be obtained by a process which comprises the steps of screening an appropriate library under stringent hybridization conditions (for example, using a temperature in the range of 45 – 65°C and an SDS concentration from 0.1 – 1%) with a labeled or detectable probe consisting of or comprising the sequence of SEQ ID NO:1 or 3 or a fragment thereof; and isolating a full-length gene and/or genomic clones containing said polynucleotide sequence.

The invention provides a polynucleotide sequence identical over its entire length to a coding sequence (open reading frame) in SEQ ID NO:1 or 3. Also provided by the invention is a coding sequence for a mature polypeptide or a fragment thereof, by itself as well as a coding sequence for a mature polypeptide or a fragment in reading frame with another coding sequence, such as a sequence encoding a leader or secretory sequence, a pre-, or pro- or prepro-protein sequence. The polynucleotide of the invention may also contain at least one non-coding sequence, including for example, but not limited to at least one non-coding 5' and 3' sequence, such as the transcribed but non-translated sequences, termination signals (such as rho-dependent and rho-independent termination signals), ribosome binding sites, Kozak sequences, sequences that stabilize mRNA, introns, and polyadenylation signals. The polynucleotide sequence may also comprise additional coding sequence encoding additional amino acids. For example, a marker sequence that facilitates purification of the fused polypeptide can be encoded. In certain embodiments of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz *et al.*, *Proc. Natl. Acad. Sci., USA* 86: 821-824 (1989), or an HA peptide tag (Wilson *et al.*, *Cell* 37: 767 (1984), both of which may be useful in purifying polypeptide sequence fused to them. Polynucleotides of the invention also include, but are not limited to, polynucleotides comprising a structural gene and its naturally associated sequences that control gene expression.

The nucleotide sequence encoding BASB112 polypeptide of SEQ ID NO:2 or 4 may be identical to the polypeptide encoding sequence contained in nucleotides 1 to 366 of SEQ ID NO:1 or the polypeptide encoding sequence contained in nucleotides 1 to 363 of SEQ ID NO:3 respectively. Alternatively it may be a sequence, which as a result of the redundancy (degeneracy) of the genetic code, also encodes the polypeptide of SEQ ID NO:2 or 4.

The term "polynucleotide encoding a polypeptide" as used herein encompasses polynucleotides that include a sequence encoding a polypeptide of the invention, particularly a bacterial polypeptide and more particularly a polypeptide of the *Moraxella catarrhalis* BASB112 having an amino acid sequence set out in SEQ ID NO:2 or 4. The term also encompasses polynucleotides that include a single continuous region or discontinuous regions encoding the polypeptide (for example, polynucleotides interrupted by integrated phage, an integrated insertion sequence, an integrated vector sequence, an integrated transposon sequence, or due to RNA editing or genomic DNA reorganization) together with additional regions, that also may contain coding and/or non-coding sequences.

The invention further relates to variants of the polynucleotides described herein that encode variants of a polypeptide having a deduced amino acid sequence of SEQ ID NO:2 or 4. Fragments of polynucleotides of the invention may be used, for example, to synthesize full-length polynucleotides of the invention.

Further particularly preferred embodiments are polynucleotides encoding BASB112 variants, that have the amino acid sequence of BASB112 polypeptide of SEQ ID NO:2 or 4 in which several, a few, 5 to 10, 1 to 5, 1 to 3, 2, 1 or no amino acid residues are substituted, modified, deleted and/or added, in any combination. Especially preferred among these are silent substitutions, additions and deletions, that do not alter the properties and activities of BASB112 polypeptide.

Further preferred embodiments of the invention are polynucleotides that are at least 85% identical over their entire length to a polynucleotide encoding BASB112 polypeptide having an amino acid sequence set out in SEQ ID NO:2 or 4, and polynucleotides that are complementary to such polynucleotides. Alternatively, most highly preferred are polynucleotides that comprise a region that is at least 90% identical over its entire length to a polynucleotide encoding BASB112 polypeptide and polynucleotides complementary thereto. In this regard, polynucleotides at least 95% identical over their entire length to the same are particularly preferred. Furthermore, those with at least 97% are highly preferred among those with at least 95%, and among these those with at least 98% and at least 99% are particularly highly preferred, with at least 99% being the more preferred.

Preferred embodiments are polynucleotides encoding polypeptides that retain substantially the same biological function or activity as the mature polypeptide encoded by a DNA of SEQ ID NO:1 or 3.

In accordance with certain preferred embodiments of this invention there are provided polynucleotides that hybridize, particularly under stringent conditions, to BASB112 polynucleotide sequences, such as those polynucleotides in SEQ ID NO:1 or 3.

The invention further relates to polynucleotides that hybridize to the polynucleotide sequences provided herein. In this regard, the invention especially relates to polynucleotides that hybridize under stringent conditions to the polynucleotides described herein. As herein used, the terms "stringent conditions" and "stringent hybridization conditions" mean hybridization occurring only if there is at least 95% and preferably at least 97% identity between the sequences. A specific example of stringent hybridization conditions is overnight incubation at 42°C in a solution comprising: 50% formamide, 5x SSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 micrograms/ml of denatured, sheared salmon sperm DNA, followed by washing the hybridization support in 0.1x SSC at about 65°C.

Hybridization and wash conditions are well known and exemplified in Sambrook, *et al.*, Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor, N.Y., (1989), particularly Chapter 11 therein. Solution hybridization may also be used with the polynucleotide sequences provided by the invention.

The invention also provides a polynucleotide consisting of or comprising a polynucleotide sequence obtained by screening an appropriate library containing the complete gene for a polynucleotide sequence set forth in SEQ ID NO:1 or 3 under stringent hybridization conditions with a probe having the sequence of said polynucleotide sequence set forth in SEQ ID NO:1 or 3 or a fragment thereof; and isolating said polynucleotide sequence. Fragments useful for obtaining such a polynucleotide include, for example, probes and primers fully described elsewhere herein.

As discussed elsewhere herein regarding polynucleotide assays of the invention, for instance, the polynucleotides of the invention, may be used as a hybridization probe for RNA, cDNA and genomic DNA to isolate full-length cDNAs and genomic clones encoding BASB112 and to isolate cDNA and genomic clones of other genes that have a high identity, particularly high sequence identity, to the BASB112 gene. Such probes generally will comprise at least 15 nucleotide residues or base pairs. Preferably, such probes will have at least 30 nucleotide residues or base pairs and may have at least 50 nucleotide residues or base pairs. Particularly preferred probes will have at least 20 nucleotide residues or base pairs and will have less than 30 nucleotide residues or base pairs.

A coding region of a BASB112 gene may be isolated by screening using a DNA sequence provided in SEQ ID NO:1 or 3 to synthesize an oligonucleotide probe. A labeled oligonucleotide having a sequence complementary to that of a gene of the invention is then used to screen a library of cDNA, genomic DNA or mRNA to determine which members of the library the probe hybridizes to.

There are several methods available and well known to those skilled in the art to obtain full-length DNAs, or extend short DNAs, for example those based on the method of Rapid Amplification of cDNA ends (RACE) (see, for example, Frohman, *et al.*, *PNAS USA* 85: 8998-9002, 1988). Recent modifications of the technique, exemplified by the Marathon™ technology (Clontech Laboratories Inc.) for example, have significantly simplified the search for longer cDNAs. In the Marathon™ technology, cDNAs have been prepared from mRNA extracted from a chosen tissue and an 'adaptor' sequence ligated onto each end. Nucleic acid amplification (PCR) is then carried out to amplify the "missing" 5' end of the DNA using a combination of gene specific and adaptor specific oligonucleotide primers. The PCR reaction is then repeated using "nested" primers, that is, primers designed to anneal within the amplified product (typically an adaptor specific primer that anneals further 3' in the adaptor sequence and a gene specific primer that anneals further 5' in the selected gene sequence). The products of this reaction can then be analyzed by DNA sequencing and a full-length DNA constructed either by joining the product directly to the existing DNA to give a complete sequence, or carrying out a separate full-length PCR using the new sequence information for the design of the 5' primer.

The polynucleotides and polypeptides of the invention may be employed, for example, as research reagents and materials for discovery of treatments of and diagnostics for diseases, particularly human diseases, as further discussed herein relating to polynucleotide assays.

The polynucleotides of the invention that are oligonucleotides derived from a sequence of SEQ ID NOS:1 or 3 may be used in the processes herein as described, but preferably for PCR, to determine whether or not the polynucleotides identified herein in whole or in part are transcribed in bacteria in infected tissue. It is recognized that such sequences will also have utility in diagnosis of the stage of infection and type of infection the pathogen has attained.

The invention also provides polynucleotides that encode a polypeptide that is the mature protein plus additional amino or carboxyl-terminal amino acids, or amino acids interior to the mature polypeptide (when the mature form has more than one polypeptide chain, for instance). Such sequences may play a role in processing of a protein from precursor to a mature form, may allow protein transport, may lengthen or shorten protein half-life or may facilitate manipulation of a protein for assay or production, among other things. As generally is the case *in vivo*, the additional amino acids may be processed away from the mature protein by cellular enzymes.

For each and every polynucleotide of the invention there is provided a polynucleotide complementary to it. It is preferred that these complementary polynucleotides are fully complementary to each polynucleotide with which they are complementary.

A precursor protein, having a mature form of the polypeptide fused to one or more prosequences may be an inactive form of the polypeptide. When prosequences are removed such inactive precursors generally are activated. Some or all of the prosequences may be removed before activation. Generally, such precursors are called proproteins.

In addition to the standard A, G, C, T/U representations for nucleotides, the term "N" may also be used in describing certain polynucleotides of the invention. "N" means that any of the four DNA or RNA nucleotides may appear at such a designated position in the DNA or RNA sequence, except it is preferred that N is not a nucleic acid that when taken in combination with adjacent nucleotide positions, when read in the correct reading frame, would have the effect of generating a premature termination codon in such reading frame.

In sum, a polynucleotide of the invention may encode a mature protein, a mature protein plus a leader sequence (which may be referred to as a preprotein), a precursor of a mature protein having one or more prosequences that are not the leader sequences of a preprotein, or a preproprotein, which is a precursor to a proprotein, having a leader sequence and one or

more prosequences, which generally are removed during processing steps that produce active and mature forms of the polypeptide.

In accordance with an aspect of the invention, there is provided the use of a polynucleotide of the invention for therapeutic or prophylactic purposes, in particular genetic immunization.

The use of a polynucleotide of the invention in genetic immunization will preferably employ a suitable delivery method such as direct injection of plasmid DNA into muscles (Wolff *et al.*, *Hum Mol Genet* (1992) 1: 363, Manthorpe *et al.*, *Hum. Gene Ther.* (1983) 4: 419), delivery of DNA complexed with specific protein carriers (Wu *et al.*, *J Biol Chem.* (1989) 264: 16985), coprecipitation of DNA with calcium phosphate (Benvenisty & Reshef, *PNAS USA*, (1986) 83: 9551), encapsulation of DNA in various forms of liposomes (Kaneda *et al.*, *Science* (1989) 243: 375), particle bombardment (Tang *et al.*, *Nature* (1992) 356:152, Eisenbraun *et al.*, *DNA Cell Biol* (1993) 12: 791) and *in vivo* infection using cloned retroviral vectors (Seeger *et al.*, *PNAS USA* (1984) 81: 5849).

Vectors, Host Cells, Expression Systems

The invention also relates to vectors that comprise a polynucleotide or polynucleotides of the invention, host cells that are genetically engineered with vectors of the invention and the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the invention.

Recombinant polypeptides of the present invention may be prepared by processes well known in those skilled in the art from genetically engineered host cells comprising expression systems. Accordingly, in a further aspect, the present invention relates to expression systems that comprise a polynucleotide or polynucleotides of the present

invention, to host cells which are genetically engineered with such expression systems, and to the production of polypeptides of the invention by recombinant techniques.

For recombinant production of the polypeptides of the invention, host cells can be genetically engineered to incorporate expression systems or portions thereof or polynucleotides of the invention. Introduction of a polynucleotide into the host cell can be effected by methods described in many standard laboratory manuals, such as Davis, *et al.*, *BASIC METHODS IN MOLECULAR BIOLOGY*, (1986) and Sambrook, *et al.*, *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989), such as, calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction and infection.

Representative examples of appropriate hosts include bacterial cells, such as cells of streptococci, staphylococci, enterococci, *E. coli*, streptomyces, cyanobacteria, *Bacillus subtilis*, *Neisseria meningitidis* and *Moraxella catarrhalis*; fungal cells, such as cells of a yeast, *Kluveromyces*, *Saccharomyces*, a basidiomycete, *Candida albicans* and *Aspergillus*; insect cells such as cells of *Drosophila* S2 and *Spodoptera* Sf9; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, 293, CV-1 and Bowes melanoma cells; and plant cells, such as cells of a gymnosperm or angiosperm.

A great variety of expression systems can be used to produce the polypeptides of the invention. Such vectors include, among others, chromosomal-, episomal- and virus-derived vectors, for example, vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses, picornaviruses, retroviruses, and alphaviruses and vectors derived from combinations thereof, such as those derived from

plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression system constructs may contain control regions that regulate as well as engender expression. Generally, any system or vector suitable to maintain, propagate or express polynucleotides and/or to express a polypeptide in a host may be used for expression in this regard. The appropriate DNA sequence may be inserted into the expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook *et al.*, *MOLECULAR CLONING, A LABORATORY MANUAL*, (*supra*).

In recombinant expression systems in eukaryotes, for secretion of a translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be incorporated into the expressed polypeptide. These signals may be endogenous to the polypeptide or they may be heterologous signals.

Polypeptides of the present invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, ion metal affinity chromatography (IMAC) is employed for purification. Well known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during intracellular synthesis, isolation and or purification.

The expression system may also be a recombinant live microorganism, such as a virus or bacterium. The gene of interest can be inserted into the genome of a live recombinant virus or bacterium. Inoculation and *in vivo* infection with this live vector will lead to *in vivo* expression of the antigen and induction of immune responses. Viruses and bacteria used for this purpose are for instance: poxviruses (e.g; vaccinia, fowlpox, canarypox), alphaviruses (Sindbis virus, Semliki Forest Virus, Venezuelan Equine Encephalitis

Virus), adenoviruses, adeno-associated virus, picornaviruses (poliovirus, rhinovirus), herpesviruses (varicella zoster virus, etc), Listeria, Salmonella, Shigella, BCG. These viruses and bacteria can be virulent, or attenuated in various ways in order to obtain live vaccines. Such live vaccines also form part of the invention.

Diagnostic, Prognostic, Serotyping and Mutation Assays

This invention is also related to the use of BASB112 polynucleotides and polypeptides of the invention for use as diagnostic reagents. Detection of BASB112 polynucleotides and/or polypeptides in a eukaryote, particularly a mammal, and especially a human, will provide a diagnostic method for diagnosis of disease, staging of disease or response of an infectious organism to drugs. Eukaryotes, particularly mammals, and especially humans, particularly those infected or suspected to be infected with an organism comprising the BASB112 gene or protein, may be detected at the nucleic acid or amino acid level by a variety of well known techniques as well as by methods provided herein.

Polypeptides and polynucleotides for prognosis, diagnosis or other analysis may be obtained from a putatively infected and/or infected individual's bodily materials. Polynucleotides from any of these sources, particularly DNA or RNA, may be used directly for detection or may be amplified enzymatically by using PCR or any other amplification technique prior to analysis. RNA, particularly mRNA, cDNA and genomic DNA may also be used in the same ways. Using amplification, characterization of the species and strain of infectious or resident organism present in an individual, may be made by an analysis of the genotype of a selected polynucleotide of the organism. Deletions and insertions can be detected by a change in size of the amplified product in comparison to a genotype of a reference sequence selected from a related organism, preferably a different species of the same genus or a different strain of the same species. Point mutations can be identified by hybridizing amplified DNA to labeled BASB112 polynucleotide sequences. Perfectly or significantly matched sequences can be distinguished from imperfectly or more significantly mismatched duplexes by DNase or RNase digestion, for DNA or RNA respectively, or by detecting

differences in melting temperatures or renaturation kinetics. Polynucleotide sequence differences may also be detected by alterations in the electrophoretic mobility of polynucleotide fragments in gels as compared to a reference sequence. This may be carried out with or without denaturing agents. Polynucleotide differences may also be detected by direct DNA or RNA sequencing. See, for example, Myers *et al.*, *Science*, 230: 1242 (1985). Sequence changes at specific locations also may be revealed by nuclease protection assays, such as RNase, V1 and S1 protection assay or a chemical cleavage method. See, for example, Cotton *et al.*, *Proc. Natl. Acad. Sci., USA*, 85: 4397-4401 (1985).

In another embodiment, an array of oligonucleotides probes comprising BASB112 nucleotide sequence or fragments thereof can be constructed to conduct efficient screening of, for example, genetic mutations, serotype, taxonomic classification or identification. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability (see, for example, Chee *et al.*, *Science*, 274: 610 (1996)).

Thus in another aspect, the present invention relates to a diagnostic kit which comprises:

- (a) a polynucleotide of the present invention, preferably the nucleotide sequence of SEQ ID NO:1 or 3, or a fragment thereof ;
- (b) a nucleotide sequence complementary to that of (a);
- (c) a polypeptide of the present invention, preferably the polypeptide of SEQ ID NO:2 or 4 or a fragment thereof; or
- (d) an antibody to a polypeptide of the present invention, preferably to the polypeptide of SEQ ID NO:2 or 4.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component. Such a kit will be of use in diagnosing a disease or susceptibility to a Disease, among others.

This invention also relates to the use of polynucleotides of the present invention as diagnostic reagents. Detection of a mutated form of a polynucleotide of the invention, preferably SEQ ID NO:1 or 3, which is associated with a disease or pathogenicity will provide a diagnostic tool that can add to, or define, a diagnosis of a disease, a prognosis of a course of disease, a determination of a stage of disease, or a susceptibility to a disease, which results from under-expression, over-expression or altered expression of the polynucleotide. Organisms, particularly infectious organisms, carrying mutations in such polynucleotide may be detected at the polynucleotide level by a variety of techniques, such as those described elsewhere herein.

Cells from an organism carrying mutations or polymorphisms (allelic variations) in a polynucleotide and/or polypeptide of the invention may also be detected at the polynucleotide or polypeptide level by a variety of techniques, to allow for serotyping, for example. For example, RT-PCR can be used to detect mutations in the RNA. It is particularly preferred to use RT-PCR in conjunction with automated detection systems, such as, for example, GeneScan. RNA, cDNA or genomic DNA may also be used for the same purpose, PCR. As an example, PCR primers complementary to a polynucleotide encoding BASB112 polypeptide can be used to identify and analyze mutations.

The invention further provides primers with 1, 2, 3 or 4 nucleotides removed from the 5' and/or the 3' end. These primers may be used for, among other things, amplifying BASB112 DNA and/or RNA isolated from a sample derived from an individual, such as a bodily material. The primers may be used to amplify a polynucleotide isolated from an infected individual, such that the polynucleotide may then be subject to various techniques for elucidation of the polynucleotide sequence. In this way, mutations in the polynucleotide sequence may be detected and used to diagnose and/or prognose the infection or its stage or course, or to serotype and/or classify the infectious agent.

The invention further provides a process for diagnosing, disease, preferably bacterial infections, more preferably infections caused by *Moraxella catarrhalis*, comprising determining from a sample derived from an individual, such as a bodily material, an increased level of expression of polynucleotide having a sequence of SEQ ID NO:1 or 3. Increased or decreased expression of a BASB112 polynucleotide can be measured using any one of the methods well known in the art for the quantitation of polynucleotides, such as, for example, amplification, PCR, RT-PCR, RNase protection, Northern blotting, spectrometry and other hybridization methods.

In addition, a diagnostic assay in accordance with the invention for detecting over-expression of BASB112 polypeptide compared to normal control tissue samples may be used to detect the presence of an infection, for example. Assay techniques that can be used to determine levels of a BASB112 polypeptide, in a sample derived from a host, such as a bodily material, are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis, antibody sandwich assays, antibody detection and ELISA assays.

The polynucleotides of the invention may be used as components of polynucleotide arrays, preferably high density arrays or grids. These high density arrays are particularly useful for diagnostic and prognostic purposes. For example, a set of spots each comprising a different gene, and further comprising a polynucleotide or polynucleotides of the invention, may be used for probing, such as using hybridization or nucleic acid amplification, using a probe obtained or derived from a bodily sample, to determine the presence of a particular polynucleotide sequence or related sequence in an individual. Such a presence may indicate the presence of a pathogen, particularly *Moraxella catarrhalis*, and may be useful in diagnosing and/or prognosing disease or a course of disease. A grid comprising a number of variants of the polynucleotide sequence of SEQ ID NO:1 or 3 are preferred. Also preferred is a grid comprising a number

of variants of a polynucleotide sequence encoding the polypeptide sequence of SEQ ID NO:2 or 4.

Antibodies

The polypeptides and polynucleotides of the invention or variants thereof, or cells expressing the same can be used as immunogens to produce antibodies immunospecific for such polypeptides or polynucleotides respectively.

In certain preferred embodiments of the invention there are provided antibodies against BASB112 polypeptides or polynucleotides.

Antibodies generated against the polypeptides or polynucleotides of the invention can be obtained by administering the polypeptides and/or polynucleotides of the invention, or epitope-bearing fragments of either or both, analogues of either or both, or cells expressing either or both, to an animal, preferably a nonhuman, using routine protocols. For preparation of monoclonal antibodies, any technique known in the art that provides antibodies produced by continuous cell line cultures can be used. Examples include various techniques, such as those in Kohler, G. and Milstein, C., *Nature* 256: 495-497 (1975); Kozbor *et al.*, *Immunology Today* 4: 72 (1983); Cole *et al.*, pg. 77-96 in *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, Alan R. Liss, Inc. (1985).

Techniques for the production of single chain antibodies (U.S. Patent No. 4,946,778) can be adapted to produce single chain antibodies to polypeptides or polynucleotides of this invention. Also, transgenic mice, or other organisms or animals, such as other mammals, may be used to express humanized antibodies immunospecific to the polypeptides or polynucleotides of the invention.

Alternatively, phage display technology may be utilized to select antibody genes with binding activities towards a polypeptide of the invention either from repertoires of PCR

amplified v-genes of lymphocytes from humans screened for possessing anti-BASB112 or from naive libraries (McCafferty, *et al.*, (1990), *Nature* 348, 552-554; Marks, *et al.*, (1992) *Biotechnology* 10, 779-783). The affinity of these antibodies can also be improved by, for example, chain shuffling (Clackson *et al.*, (1991) *Nature* 352: 628).

The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptides or polynucleotides of the invention to purify the polypeptides or polynucleotides by, for example, affinity chromatography.

Thus, among others, antibodies against BASB112-polypeptide or BASB112-polynucleotide may be employed to treat infections, particularly bacterial infections.

Polypeptide variants include antigenically, epitopically or immunologically equivalent variants form a particular aspect of this invention.

Preferably, the antibody or variant thereof is modified to make it less immunogenic in the individual. For example, if the individual is human the antibody may most preferably be "humanized," where the complementarity determining region or regions of the hybridoma-derived antibody has been transplanted into a human monoclonal antibody, for example as described in Jones *et al.* (1986), *Nature* 321, 522-525 or Tempest *et al.*, (1991) *Biotechnology* 9, 266-273.

Antagonists and Agonists - Assays and Molecules

Polypeptides and polynucleotides of the invention may also be used to assess the binding of small molecule substrates and ligands in, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. These substrates and ligands may be natural substrates and ligands or may be structural or functional mimetics. See, *e.g.*, Coligan *et al.*, *Current Protocols in Immunology* 1(2): Chapter 5 (1991).

The screening methods may simply measure the binding of a candidate compound to the polypeptide or polynucleotide, or to cells or membranes bearing the polypeptide or polynucleotide, or a fusion protein of the polypeptide by means of a label directly or indirectly associated with the candidate compound. Alternatively, the screening method may involve competition with a labeled competitor. Further, these screening methods may test whether the candidate compound results in a signal generated by activation or inhibition of the polypeptide or polynucleotide, using detection systems appropriate to the cells comprising the polypeptide or polynucleotide. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed. Constitutively active polypeptide and/or constitutively expressed polypeptides and polynucleotides may be employed in screening methods for inverse agonists or inhibitors, in the absence of an agonist or inhibitor, by testing whether the candidate compound results in inhibition of activation of the polypeptide or polynucleotide, as the case may be. Further, the screening methods may simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide or polynucleotide of the present invention, to form a mixture, measuring BASB112 polypeptide and/or polynucleotide activity in the mixture, and comparing the BASB112 polypeptide and/or polynucleotide activity of the mixture to a standard. Fusion proteins, such as those made from Fc portion and BASB112 polypeptide, as hereinbefore described, can also be used for high-throughput screening assays to identify antagonists of the polypeptide of the present invention, as well as of phylogenetically and and/or functionally related polypeptides (see D. Bennett *et al.*, J Mol Recognition, 8:52-58 (1995); and K. Johanson *et al.*, J Biol Chem, 270(16):9459-9471 (1995)).

The polynucleotides, polypeptides and antibodies that bind to and/or interact with a polypeptide of the present invention may also be used to configure screening methods for detecting the effect of added compounds on the production of mRNA and/or polypeptide

in cells. For example, an ELISA assay may be constructed for measuring secreted or cell associated levels of polypeptide using monoclonal and polyclonal antibodies by standard methods known in the art. This can be used to discover agents which may inhibit or enhance the production of polypeptide (also called antagonist or agonist, respectively) from suitably manipulated cells or tissues.

The invention also provides a method of screening compounds to identify those which enhance (agonist) or block (antagonist) the action of BASB112 polypeptides or polynucleotides, particularly those compounds that are bacteriostatic and/or bactericidal. The method of screening may involve high-throughput techniques. For example, to screen for agonists or antagonists, a synthetic reaction mix, a cellular compartment, such as a membrane, cell envelope or cell wall, or a preparation of any thereof, comprising BASB112 polypeptide and a labeled substrate or ligand of such polypeptide is incubated in the absence or the presence of a candidate molecule that may be a BASB112 agonist or antagonist. The ability of the candidate molecule to agonize or antagonize the BASB112 polypeptide is reflected in decreased binding of the labeled ligand or decreased production of product from such substrate. Molecules that bind gratuitously, *i.e.*, without inducing the effects of BASB112 polypeptide are most likely to be good antagonists. Molecules that bind well and, as the case may be, increase the rate of product production from substrate, increase signal transduction, or increase chemical channel activity are agonists. Detection of the rate or level of, as the case may be, production of product from substrate, signal transduction, or chemical channel activity may be enhanced by using a reporter system. Reporter systems that may be useful in this regard include but are not limited to colorimetric, labeled substrate converted into product, a reporter gene that is responsive to changes in BASB112 polynucleotide or polypeptide activity, and binding assays known in the art.

Another example of an assay for BASB112 agonists is a competitive assay that combines BASB112 and a potential agonist with BASB112-binding molecules, recombinant BASB112 binding molecules, natural substrates or ligands, or substrate or ligand mimetics,

under appropriate conditions for a competitive inhibition assay. BASB112 can be labeled, such as by radioactivity or a colorimetric compound, such that the number of BASB112 molecules bound to a binding molecule or converted to product can be determined accurately to assess the effectiveness of the potential antagonist.

Potential antagonists include, among others, small organic molecules, peptides, polypeptides and antibodies that bind to a polynucleotide and/or polypeptide of the invention and thereby inhibit or extinguish its activity or expression. Potential antagonists also may be small organic molecules, a peptide, a polypeptide such as a closely related protein or antibody that binds the same sites on a binding molecule, such as a binding molecule, without inducing BASB112-induced activities, thereby preventing the action or expression of BASB112 polypeptides and/or polynucleotides by excluding BASB112 polypeptides and/or polynucleotides from binding.

Potential antagonists include a small molecule that binds to and occupies the binding site of the polypeptide thereby preventing binding to cellular binding molecules, such that normal biological activity is prevented. Examples of small molecules include but are not limited to small organic molecules, peptides or peptide-like molecules. Other potential antagonists include antisense molecules (see Okano, *J. Neurochem.* 56: 560 (1991); *OLIGODEOXYNUCLEOTIDES AS ANTISENSE INHIBITORS OF GENE EXPRESSION*, CRC Press, Boca Raton, FL (1988), for a description of these molecules). Preferred potential antagonists include compounds related to and variants of BASB112.

In a further aspect, the present invention relates to genetically engineered soluble fusion proteins comprising a polypeptide of the present invention, or a fragment thereof, and various portions of the constant regions of heavy or light chains of immunoglobulins of various subclasses (IgG, IgM, IgA, IgE). Preferred as an immunoglobulin is the constant part of the heavy chain of human IgG, particularly IgG1, where fusion takes place at the hinge region. In a particular embodiment, the Fc part can be removed simply by

incorporation of a cleavage sequence which can be cleaved with blood clotting factor Xa. Furthermore, this invention relates to processes for the preparation of these fusion proteins by genetic engineering, and to the use thereof for drug screening, diagnosis and therapy. A further aspect of the invention also relates to polynucleotides encoding such fusion proteins. Examples of fusion protein technology can be found in International Patent Application Nos. WO94/29458 and WO94/22914.

Each of the polynucleotide sequences provided herein may be used in the discovery and development of antibacterial compounds. The encoded protein, upon expression, can be used as a target for the screening of antibacterial drugs. Additionally, the polynucleotide sequences encoding the amino terminal regions of the encoded protein or Shine-Delgarno or other translation facilitating sequences of the respective mRNA can be used to construct antisense sequences to control the expression of the coding sequence of interest.

The invention also provides the use of the polypeptide, polynucleotide, agonist or antagonist of the invention to interfere with the initial physical interaction between a pathogen or pathogens and a eukaryotic, preferably mammalian, host responsible for sequelae of infection. In particular, the molecules of the invention may be used: in the prevention of adhesion of bacteria, in particular gram positive and/or gram negative bacteria, to eukaryotic, preferably mammalian, extracellular matrix proteins on in-dwelling devices or to extracellular matrix proteins in wounds; to block bacterial adhesion between eukaryotic, preferably mammalian, extracellular matrix proteins and bacterial BASB112 proteins that mediate tissue damage and/or; to block the normal progression of pathogenesis in infections initiated other than by the implantation of in-dwelling devices or by other surgical techniques.

In accordance with yet another aspect of the invention, there are provided BASB112 agonists and antagonists, preferably bacteristatic or bactericidal agonists and antagonists.

The antagonists and agonists of the invention may be employed, for instance, to prevent, inhibit and/or treat diseases.

In a further aspect, the present invention relates to mimotopes of the polypeptide of the invention. A mimotope is a peptide sequence, sufficiently similar to the native peptide (sequentially or structurally), which is capable of being recognised by antibodies which recognise the native peptide; or is capable of raising antibodies which recognise the native peptide when coupled to a suitable carrier.

Peptide mimotopes may be designed for a particular purpose by addition, deletion or substitution of elected amino acids. Thus, the peptides may be modified for the purposes of ease of conjugation to a protein carrier. For example, it may be desirable for some chemical conjugation methods to include a terminal cysteine. In addition it may be desirable for peptides conjugated to a protein carrier to include a hydrophobic terminus distal from the conjugated terminus of the peptide, such that the free unconjugated end of the peptide remains associated with the surface of the carrier protein. Thereby presenting the peptide in a conformation which most closely resembles that of the peptide as found in the context of the whole native molecule. For example, the peptides may be altered to have an N-terminal cysteine and a C-terminal hydrophobic amidated tail. Alternatively, the addition or substitution of a D-stereoisomer form of one or more of the amino acids may be performed to create a beneficial derivative, for example to enhance stability of the peptide.

Alternatively, peptide mimotopes may be identified using antibodies which are capable themselves of binding to the polypeptides of the present invention using techniques such as phage display technology (EP 0 552 267 B1). This technique, generates a large number of peptide sequences which mimic the structure of the native peptides and are, therefore, capable of binding to anti-native peptide antibodies, but may not necessarily themselves share significant sequence homology to the native polypeptide.

Vaccines

Another aspect of the invention relates to a method for inducing an immunological response in an individual, particularly a mammal, preferably humans, which comprises inoculating the individual with BASB112 polynucleotide and/or polypeptide, or a fragment or variant thereof, adequate to produce antibody and/ or T cell immune response to protect said individual from infection, particularly bacterial infection and most particularly *Moraxella catarrhalis* infection. Also provided are methods whereby such immunological response slows bacterial replication. Yet another aspect of the invention relates to a method of inducing immunological response in an individual which comprises delivering to such individual a nucleic acid vector, sequence or ribozyme to direct expression of BASB112 polynucleotide and/or polypeptide, or a fragment or a variant thereof, for expressing BASB112 polynucleotide and/or polypeptide, or a fragment or a variant thereof *in vivo* in order to induce an immunological response, such as, to produce antibody and/ or T cell immune response, including, for example, cytokine-producing T cells or cytotoxic T cells, to protect said individual, preferably a human, from disease, whether that disease is already established within the individual or not. One example of administering the gene is by accelerating it into the desired cells as a coating on particles or otherwise. Such nucleic acid vector may comprise DNA, RNA, a ribozyme, a modified nucleic acid, a DNA/RNA hybrid, a DNA-protein complex or an RNA-protein complex.

A further aspect of the invention relates to an immunological composition that when introduced into an individual, preferably a human, capable of having induced within it an immunological response, induces an immunological response in such individual to a BASB112 polynucleotide and/or polypeptide encoded therefrom, wherein the composition comprises a recombinant BASB112 polynucleotide and/or polypeptide encoded therefrom and/or comprises DNA and/or RNA which encodes and expresses an antigen of said BASB112 polynucleotide, polypeptide encoded therefrom, or other polypeptide of the invention. The immunological response may be used therapeutically or prophylactically

and may take the form of antibody immunity and/or cellular immunity, such as cellular immunity arising from CTL or CD4+ T cells.

A BASB112 polypeptide or a fragment thereof may be fused with co-protein or chemical moiety which may or may not by itself produce antibodies, but which is capable of stabilizing the first protein and producing a fused or modified protein which will have antigenic and/or immunogenic properties, and preferably protective properties. Thus fused recombinant protein, preferably further comprises an antigenic co-protein, such as lipoprotein D from *Haemophilus influenzae*, Glutathione-S-transferase (GST) or beta-galactosidase, or any other relatively large co-protein which solubilizes the protein and facilitates production and purification thereof. Moreover, the co-protein may act as an adjuvant in the sense of providing a generalized stimulation of the immune system of the organism receiving the protein. The co-protein may be attached to either the amino- or carboxy-terminus of the first protein.

In a vaccine composition according to the invention, a BASB112 polypeptide and/or polynucleotide, or a fragment, or a mimotope, or a variant thereof may be present in a vector, such as the live recombinant vectors described above for example live bacterial vectors.

Also suitable are non-live vectors for the BASB112 polypeptide, for example bacterial outer-membrane vesicles or "blebs". OM blebs are derived from the outer membrane of the two-layer membrane of Gram-negative bacteria and have been documented in many Gram-negative bacteria (Zhou, L *et al.* 1998. *FEMS Microbiol. Lett.* 163:223-228) including *C. trachomatis* and *C. psittaci*. A non-exhaustive list of bacterial pathogens reported to produce blebs also includes: *Bordetella pertussis*, *Borrelia burgdorferi*, *Brucella melitensis*, *Brucella ovis*, *Escherichia coli*, *Haemophilus influenza*, *Legionella pneumophila*, *Moraxella catarrhalis*, *Neisseria gonorrhoeae*, *Neisseria meningitidis*, *Pseudomonas aeruginosa* and *Yersinia enterocolitica*.

Blebs have the advantage of providing outer-membrane proteins in their native conformation and are thus particularly useful for vaccines. Blebs can also be improved for vaccine use by engineering the bacterium so as to modify the expression of one or more molecules at the outer membrane. Thus for example the expression of a desired immunogenic protein at the outer membrane, such as the BASB112 polypeptide, can be introduced or upregulated (e.g. by altering the promoter). Instead or in addition, the expression of outer-membrane molecules which are either not relevant (e.g. unprotective antigens or immunodominant but variable proteins) or detrimental (e.g. toxic molecules such as LPS, or potential inducers of an autoimmune response) can be downregulated. These approaches are discussed in more detail below.

The non-coding flanking regions of the BASB112 gene contain regulatory elements important in the expression of the gene. This regulation takes place both at the transcriptional and translational level. The sequence of these regions, either upstream or downstream of the open reading frame of the gene, can be obtained by DNA sequencing. This sequence information allows the determination of potential regulatory motifs such as the different promoter elements, terminator sequences, inducible sequence elements, repressors, elements responsible for phase variation, the shine-dalgarno sequence, regions with potential secondary structure involved in regulation, as well as other types of regulatory motifs or sequences. This sequence is a further aspect of the invention.

This sequence information allows the modulation of the natural expression of the BASB112 gene. The upregulation of the gene expression may be accomplished by altering the promoter, the shine-dalgarno sequence, potential repressor or operator elements, or any other elements involved. Likewise, downregulation of expression can be achieved by similar types of modification. Alternatively, by changing phase variation sequences, the expression of the gene can be put under phase variation control, or it may be uncoupled from this regulation. In another approach, the expression of the gene can be

put under the control of one or more inducible elements allowing regulated expression. Examples of such regulation include, but are not limited to, induction by temperature shift, addition of inductor substrates like selected carbohydrates or their derivatives, trace elements, vitamins, co-factors, metal ions, etc.

Such modifications as described above can be introduced by several different means. The modification of sequences involved in gene expression can be carried out *in vivo* by random mutagenesis followed by selection for the desired phenotype. Another approach consists in isolating the region of interest and modifying it by random mutagenesis, or site-directed replacement, insertion or deletion mutagenesis. The modified region can then be reintroduced into the bacterial genome by homologous recombination, and the effect on gene expression can be assessed. In another approach, the sequence knowledge of the region of interest can be used to replace or delete all or part of the natural regulatory sequences. In this case, the regulatory region targeted is isolated and modified so as to contain the regulatory elements from another gene, a combination of regulatory elements from different genes, a synthetic regulatory region, or any other regulatory region, or to delete selected parts of the wild-type regulatory sequences. These modified sequences can then be reintroduced into the bacterium via homologous recombination into the genome. A non-exhaustive list of preferred promoters that could be used for up-regulation of gene expression includes the promoters *porA*, *porB*, *lbpB*, *tbpB*, *p112*, *lst*, *hpuAB* from *N. meningitidis* or *N. gonorrhoeae*; *ompCD*, *copB*, *lbpB*, *ompE*, *UspA1*; *UspA2*; *TbpB* from *M. Catarrhalis*; *p1*, *p2*, *p4*, *p5*, *p6*, *lpD*, *tbpB*, *D15*, *Hia*, *Hmw1*, *Hmw2* from *H. influenzae*.

In one example, the expression of the gene can be modulated by exchanging its promoter with a stronger promoter (through isolating the upstream sequence of the gene, *in vitro* modification of this sequence, and reintroduction into the genome by homologous recombination). Upregulated expression can be obtained in both the bacterium as well as in the outer membrane vesicles shed (or made) from the bacterium.

In other examples, the described approaches can be used to generate recombinant bacterial strains with improved characteristics for vaccine applications. These can be, but are not limited to, attenuated strains, strains with increased expression of selected antigens, strains with knock-outs (or decreased expression) of genes interfering with the immune response, strains with modulated expression of immunodominant proteins, strains with modulated shedding of outer-membrane vesicles.

Thus, also provided by the invention is a modified upstream region of the BASB112 gene, which modified upstream region contains a heterologous regulatory element which alters the expression level of the BASB112 protein located at the outer membrane. The upstream region according to this aspect of the invention includes the sequence upstream of the BASB112 gene. The upstream region starts immediately upstream of the BASB112 gene and continues usually to a position no more than about 1000 bp upstream of the gene from the ATG start codon. In the case of a gene located in a polycistronic sequence (operon) the upstream region can start immediately preceding the gene of interest, or preceding the first gene in the operon. Preferably, a modified upstream region according to this aspect of the invention contains a heterologous promotor at a position between 500 and 700 bp upstream of the ATG.

Thus, the invention provides a BASB112 polypeptide, in a modified bacterial bleb. The invention further provides modified host cells capable of producing the non-live membrane-based bleb vectors. The invention further provides nucleic acid vectors comprising the BASB112 gene having a modified upstream region containing a heterologous regulatory element.

Further provided by the invention are processes to prepare the host cells and bacterial blebs according to the invention.

Provided by this invention are compositions, particularly vaccine compositions, and methods comprising the polypeptides and/or polynucleotides of the invention and immunostimulatory DNA sequences, such as those described in Sato, Y. *et al.* Science 273: 352 (1996).

Also, provided by this invention are methods using the described polynucleotide or particular fragments thereof, which have been shown to encode non-variable regions of bacterial cell surface proteins, in polynucleotide constructs used in such genetic immunization experiments in animal models of infection with *Moraxella catarrhalis*. Such experiments will be particularly useful for identifying protein epitopes able to provoke a prophylactic or therapeutic immune response. It is believed that this approach will allow for the subsequent preparation of monoclonal antibodies of particular value, derived from the requisite organ of the animal successfully resisting or clearing infection, for the development of prophylactic agents or therapeutic treatments of bacterial infection, particularly *Moraxella catarrhalis* infection, in mammals, particularly humans.

The invention also includes a vaccine formulation which comprises an immunogenic recombinant polypeptide and/or polynucleotide of the invention together with a suitable carrier, such as a pharmaceutically acceptable carrier. Since the polypeptides and polynucleotides may be broken down in the stomach, each is preferably administered parenterally, including, for example, administration that is subcutaneous, intramuscular, intravenous, or intradermal. Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostatic compounds and solutes which render the formulation isotonic with the bodily fluid, preferably the blood, of the individual; and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use.

The vaccine formulation of the invention may also include adjuvant systems for enhancing the immunogenicity of the formulation. Preferably the adjuvant system raises preferentially a TH1 type of response.

An immune response may be broadly distinguished into two extreme categories, being a humoral or cell mediated immune responses (traditionally characterised by antibody and cellular effector mechanisms of protection respectively). These categories of response have been termed TH1-type responses (cell-mediated response), and TH2-type immune responses (humoral response).

Extreme TH1-type immune responses may be characterised by the generation of antigen specific, haplotype restricted cytotoxic T lymphocytes, and natural killer cell responses. In mice TH1-type responses are often characterised by the generation of antibodies of the IgG2a subtype, whilst in the human these correspond to IgG1 type antibodies. TH2-type immune responses are characterised by the generation of a broad range of immunoglobulin isotypes including in mice IgG1, IgA, and IgM.

It can be considered that the driving force behind the development of these two types of immune responses are cytokines. High levels of TH1-type cytokines tend to favour the induction of cell mediated immune responses to the given antigen, whilst high levels of TH2-type cytokines tend to favour the induction of humoral immune responses to the antigen.

The distinction of TH1 and TH2-type immune responses is not absolute. In reality an individual will support an immune response which is described as being predominantly TH1 or predominantly TH2. However, it is often convenient to consider the families of cytokines in terms of that described in murine CD4 +ve T cell clones by Mosmann and Coffman (*Mosmann, T.R. and Coffman, R.L. (1989) TH1 and TH2 cells: different*

patterns of lymphokine secretion lead to different functional properties. Annual Review of Immunology, 7, p145-173). Traditionally, TH1-type responses are associated with the production of the INF- γ and IL-2 cytokines by T-lymphocytes. Other cytokines often directly associated with the induction of TH1-type immune responses are not produced by T-cells, such as IL-12. In contrast, TH2- type responses are associated with the secretion of IL-4, IL-5, IL-6 and IL-13.

It is known that certain vaccine adjuvants are particularly suited to the stimulation of either TH1 or TH2 - type cytokine responses. Traditionally the best indicators of the TH1:TH2 balance of the immune response after a vaccination or infection includes direct measurement of the production of TH1 or TH2 cytokines by T lymphocytes *in vitro* after restimulation with antigen, and/or the measurement of the IgG1:IgG2a ratio of antigen specific antibody responses.

Thus, a TH1-type adjuvant is one which preferentially stimulates isolated T-cell populations to produce high levels of TH1-type cytokines when re-stimulated with antigen *in vitro*, and promotes development of both CD8+ cytotoxic T lymphocytes and antigen specific immunoglobulin responses associated with TH1-type isotype.

Adjuvants which are capable of preferential stimulation of the TH1 cell response are described in International Patent Application No. WO 94/00153 and WO 95/17209.

3 De-O-acylated monophosphoryl lipid A (3D-MPL) is one such adjuvant. This is known from GB 2220211 (Ribi). Chemically it is a mixture of 3 De-O-acylated monophosphoryl lipid A with 4, 5 or 6 acylated chains and is manufactured by Ribi Immunochem, Montana. A preferred form of 3 De-O-acylated monophosphoryl lipid A is disclosed in European Patent 0 689 454 B1 (SmithKline Beecham Biologicals SA).

Preferably, the particles of 3D-MPL are small enough to be sterile filtered through a 0.22micron membrane (European Patent number 0 689 454).

3D-MPL will be present in the range of 10 μ g - 100 μ g preferably 25-50 μ g per dose wherein the antigen will typically be present in a range 2-50 μ g per dose.

Another preferred adjuvant comprises QS21, an Hplc purified non-toxic fraction derived from the bark of Quillaja Saponaria Molina. Optionally this may be admixed with 3 De-O-acylated monophosphoryl lipid A (3D-MPL), optionally together with an carrier.

The method of production of QS21 is disclosed in US patent No. 5,057,540.

Non-reactogenic adjuvant formulations containing QS21 have been described previously (WO 96/33739). Such formulations comprising QS21 and cholesterol have been shown to be successful TH1 stimulating adjuvants when formulated together with an antigen.

Further adjuvants which are preferential stimulators of TH1 cell response include immunomodulatory oligonucleotides, for example unmethylated CpG sequences as disclosed in WO 96/02555.

Combinations of different TH1 stimulating adjuvants, such as those mentioned hereinabove, are also contemplated as providing an adjuvant which is a preferential stimulator of TH1 cell response. For example, QS21 can be formulated together with 3D-MPL. The ratio of QS21 : 3D-MPL will typically be in the order of 1 : 10 to 10 : 1; preferably 1:5 to 5 : 1 and often substantially 1 : 1. The preferred range for optimal synergy is 2.5 : 1 to 1 : 1 3D-MPL: QS21.

Preferably a carrier is also present in the vaccine composition according to the invention. The carrier may be an oil in water emulsion, or an aluminium salt, such as aluminium phosphate or aluminium hydroxide.

A preferred oil-in-water emulsion comprises a metabolisable oil, such as squalene, alpha tocopherol and Tween 80. In a particularly preferred aspect the antigens in the vaccine composition according to the invention are combined with QS21 and 3D-MPL in such an emulsion. Additionally the oil in water emulsion may contain span 85 and/or lecithin and/or tricaprylin.

Typically for human administration QS21 and 3D-MPL will be present in a vaccine in the range of 1µg - 200µg, such as 10-100µg, preferably 10µg - 50µg per dose.

Typically the oil in water will comprise from 2 to 10% squalene, from 2 to 10% alpha tocopherol and from 0.3 to 3% tween 80. Preferably the ratio of squalene: alpha tocopherol is equal to or less than 1 as this provides a more stable emulsion. Span 85 may also be present at a level of 1%. In some cases it may be advantageous that the vaccines of the present invention will further contain a stabiliser.

Non-toxic oil in water emulsions preferably contain a non-toxic oil, e.g. squalane or squalene, an emulsifier, e.g. Tween 80, in an aqueous carrier. The aqueous carrier may be, for example, phosphate buffered saline.

A particularly potent adjuvant formulation involving QS21, 3D-MPL and tocopherol in an oil in water emulsion is described in WO 95/17210.

The present invention also provides a polyvalent vaccine composition comprising a vaccine formulation of the invention in combination with other antigens, in particular antigens useful for treating cancers, autoimmune diseases and related conditions. Such a

polyvalent vaccine composition may include a TH-1 inducing adjuvant as hereinbefore described.

While the invention has been described with reference to certain BASB112 polypeptides and polynucleotides, it is to be understood that this covers fragments of the naturally occurring polypeptides and polynucleotides, and similar polypeptides and polynucleotides with additions, deletions or substitutions which do not substantially affect the immunogenic properties of the recombinant polypeptides or polynucleotides.

Compositions, kits and administration

In a further aspect of the invention there are provided compositions comprising a BASB112 polynucleotide and/or a BASB112 polypeptide for administration to a cell or to a multicellular organism.

The invention also relates to compositions comprising a polynucleotide and/or a polypeptides discussed herein or their agonists or antagonists. The polypeptides and polynucleotides of the invention may be employed in combination with a non-sterile or sterile carrier or carriers for use with cells, tissues or organisms, such as a pharmaceutical carrier suitable for administration to an individual. Such compositions comprise, for instance, a media additive or a therapeutically effective amount of a polypeptide and/or polynucleotide of the invention and a pharmaceutically acceptable carrier or excipient. Such carriers may include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol and combinations thereof. The formulation should suit the mode of administration. The invention further relates to diagnostic and pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

Polypeptides, polynucleotides and other compounds of the invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

The pharmaceutical compositions may be administered in any effective, convenient manner including, for instance, administration by topical, oral, anal, vaginal, intravenous, intraperitoneal, intramuscular, subcutaneous, intranasal or intradermal routes among others.

In therapy or as a prophylactic, the active agent may be administered to an individual as an injectable composition, for example as a sterile aqueous dispersion, preferably isotonic.

In a further aspect, the present invention provides for pharmaceutical compositions comprising a therapeutically effective amount of a polypeptide and/or polynucleotide, such as the soluble form of a polypeptide and/or polynucleotide of the present invention, agonist or antagonist peptide or small molecule compound, in combination with a pharmaceutically acceptable carrier or excipient. Such carriers include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. The invention further relates to pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention.

Polypeptides, polynucleotides and other compounds of the present invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

The composition will be adapted to the route of administration, for instance by a systemic or an oral route. Preferred forms of systemic administration include injection, typically by intravenous injection. Other injection routes, such as subcutaneous, intramuscular, or intraperitoneal, can be used. Alternative means for systemic administration include transmucosal and transdermal administration using penetrants such as bile salts or fusidic acids or other detergents. In addition, if a polypeptide or other compounds of the present invention can be formulated in an enteric or an encapsulated formulation, oral administration may also be possible. Administration of these compounds may also be topical and/or localized, in the form of salves, pastes, gels, solutions, powders and the like.

For administration to mammals, and particularly humans, it is expected that the daily dosage level of the active agent will be from 0.01 mg/kg to 10 mg/kg, typically around 1 mg/kg. The physician in any event will determine the actual dosage which will be most suitable for an individual and will vary with the age, weight and response of the particular individual. The above dosages are exemplary of the average case. There can, of course, be individual instances where higher or lower dosage ranges are merited, and such are within the scope of this invention.

The dosage range required depends on the choice of peptide, the route of administration, the nature of the formulation, the nature of the subject's condition, and the judgment of the attending practitioner. Suitable dosages, however, are in the range of 0.1-100 μ g/kg of subject.

A vaccine composition is conveniently in injectable form. Conventional adjuvants may be employed to enhance the immune response. A suitable unit dose for vaccination is 0.5-5 microgram/kg of antigen, and such dose is preferably administered 1-3 times and with an interval of 1-3 weeks. With the indicated dose range, no adverse toxicological effects will be observed with the compounds of the invention which would preclude their administration to suitable individuals.

Wide variations in the needed dosage, however, are to be expected in view of the variety of compounds available and the differing efficiencies of various routes of administration. For example, oral administration would be expected to require higher dosages than administration by intravenous injection. Variations in these dosage levels can be adjusted using standard empirical routines for optimization, as is well understood in the art.

Sequence Databases, Sequences in a Tangible Medium, and Algorithms

Polynucleotide and polypeptide sequences form a valuable information resource with which to determine their 2- and 3-dimensional structures as well as to identify further sequences of similar homology. These approaches are most easily facilitated by storing the sequence in a computer readable medium and then using the stored data in a known macromolecular structure program or to search a sequence database using well known searching tools, such as the GCG program package.

Also provided by the invention are methods for the analysis of character sequences or strings, particularly genetic sequences or encoded protein sequences. Preferred methods of sequence analysis include, for example, methods of sequence homology analysis, such as identity and similarity analysis, DNA, RNA and protein structure analysis, sequence assembly, cladistic analysis, sequence motif analysis, open reading frame determination, nucleic acid base calling, codon usage analysis, nucleic acid base trimming, and sequencing chromatogram peak analysis.

A computer based method is provided for performing homology identification. This method comprises the steps of: providing a first polynucleotide sequence comprising the sequence of a polynucleotide of the invention in a computer readable medium; and comparing said first polynucleotide sequence to at least one second polynucleotide or polypeptide sequence to identify homology.

A computer based method is also provided for performing homology identification, said method comprising the steps of: providing a first polypeptide sequence comprising the sequence of a polypeptide of the invention in a computer readable medium; and comparing said first polypeptide sequence to at least one second polynucleotide or polypeptide sequence to identify homology.

All publications and references, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference in their

entirety as if each individual publication or reference were specifically and individually indicated to be incorporated by reference herein as being fully set forth. Any patent application to which this application claims priority is also incorporated by reference herein in its entirety in the manner described above for publications and references.

DEFINITIONS

"Identity," as known in the art, is a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, as the case may be, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. "Identity" can be readily calculated by known methods, including but not limited to those described in (*Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data*, Part I, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heine, G., Academic Press, 1987; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; and Carillo, H., and Lipman, D., *SIAM J. Applied Math.*, 48: 1073 (1988). Methods to determine identity are designed to give the largest match between the sequences tested. Moreover, methods to determine identity are codified in publicly available computer programs. Computer program methods to determine identity between two sequences include, but are not limited to, the GAP program in the GCG program package (Devereux, J., et al., *Nucleic Acids Research* 12(1): 387 (1984)), BLASTP, BLASTN (Altschul, S.F. et al., *J. Molec. Biol.* 215: 403-410 (1990), and FASTA(Pearson and Lipman *Proc. Natl. Acad. Sci. USA* 85: 2444-2448 (1988). The BLAST family of programs is publicly available from NCBI and other sources (*BLAST Manual*, Altschul, S., et al., NCBI NLM NIH Bethesda, MD 20894;

Altschul, S., *et al.*, *J. Mol. Biol.* 215: 403-410 (1990). The well known Smith Waterman algorithm may also be used to determine identity.

Parameters for polypeptide sequence comparison include the following:

Algorithm: Needleman and Wunsch, *J. Mol Biol.* 48: 443-453 (1970)

Comparison matrix: BLOSSUM62 from Henikoff and Henikoff,
Proc. Natl. Acad. Sci. USA. 89:10915-10919 (1992)

Gap Penalty: 8

Gap Length Penalty: 2

A program useful with these parameters is publicly available as the "gap" program from Genetics Computer Group, Madison WI. The aforementioned parameters are the default parameters for peptide comparisons (along with no penalty for end gaps).

Parameters for polynucleotide comparison include the following:

Algorithm: Needleman and Wunsch, *J. Mol Biol.* 48: 443-453 (1970)

Comparison matrix: matches = +10, mismatch = 0

Gap Penalty: 50

Gap Length Penalty: 3

Available as: The "gap" program from Genetics Computer Group, Madison WI. These are the default parameters for nucleic acid comparisons.

A preferred meaning for "identity" for polynucleotides and polypeptides, as the case may be, are provided in (1) and (2) below.

(1) Polynucleotide embodiments further include an isolated polynucleotide comprising a polynucleotide sequence having at least a 50, 60, 70, 80, 85, 90, 95, 97 or 100% identity to the reference sequence of SEQ ID NO:1, wherein said polynucleotide sequence may be identical to the reference sequence of SEQ ID NO:1 or may include up to a certain integer number of nucleotide alterations as compared to the reference

sequence, wherein said alterations are selected from the group consisting of at least one nucleotide deletion, substitution, including transition and transversion, or insertion, and wherein said alterations may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among the nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence, and wherein said number of nucleotide alterations is determined by multiplying the total number of nucleotides in SEQ ID NO:1 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of nucleotides in SEQ ID NO:1, or:

$$n_n \leq x_n - (x_n \circ y),$$

wherein n_n is the number of nucleotide alterations, x_n is the total number of nucleotides in SEQ ID NO:1, y is 0.50 for 50%, 0.60 for 60%, 0.70 for 70%, 0.80 for 80%, 0.85 for 85%, 0.90 for 90%, 0.95 for 95%, 0.97 for 97% or 1.00 for 100%, and \circ is the symbol for the multiplication operator, and wherein any non-integer product of x_n and y is rounded down to the nearest integer prior to subtracting it from x_n . Alterations of a polynucleotide sequence encoding the polypeptide of SEQ ID NO:2 may create nonsense, missense or frameshift mutations in this coding sequence and thereby alter the polypeptide encoded by the polynucleotide following such alterations.

By way of example, a polynucleotide sequence of the present invention may be identical to the reference sequence of SEQ ID NO:1, that is it may be 100% identical, or it may include up to a certain integer number of nucleic acid alterations as compared to the reference sequence such that the percent identity is less than 100% identity. Such alterations are selected from the group consisting of at least one nucleic acid deletion, substitution, including transition and transversion, or insertion, and wherein said alterations may occur at the 5' or 3' terminal positions of the reference polynucleotide sequence or anywhere between those terminal positions, interspersed either individually

among the nucleic acids in the reference sequence or in one or more contiguous groups within the reference sequence. The number of nucleic acid alterations for a given percent identity is determined by multiplying the total number of nucleic acids in SEQ ID NO:1 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of nucleic acids in SEQ ID NO:1, or:

$$n_n \leq x_n - (x_n \circ y),$$

wherein n_n is the number of nucleic acid alterations, x_n is the total number of nucleic acids in SEQ ID NO:1, y is, for instance 0.70 for 70%, 0.80 for 80%, 0.85 for 85% etc., \circ is the symbol for the multiplication operator, and wherein any non-integer product of x_n and y is rounded down to the nearest integer prior to subtracting it from x_n .

(2) Polypeptide embodiments further include an isolated polypeptide comprising a polypeptide having at least a 50,60, 70, 80, 85, 90, 95, 97 or 100% identity to a polypeptide reference sequence of SEQ ID NO:2, wherein said polypeptide sequence may be identical to the reference sequence of SEQ ID NO:2 or may include up to a certain integer number of amino acid alterations as compared to the reference sequence, wherein said alterations are selected from the group consisting of at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion, and wherein said alterations may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between those terminal positions, interspersed either individually among the amino acids in the reference sequence or in one or more contiguous groups within the reference sequence, and wherein said number of amino acid alterations is determined by multiplying the total number of amino acids in SEQ ID NO:2 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of amino acids in SEQ ID NO:2, or:

$$n_a \leq x_a - (x_a \circ y),$$

wherein n_a is the number of amino acid alterations, x_a is the total number of amino acids in SEQ ID NO:2, y is 0.50 for 50%, 0.60 for 60%, 0.70 for 70%, 0.80 for 80%, 0.85 for 85%, 0.90 for 90%, 0.95 for 95%, 0.97 for 97% or 1.00 for 100%, and \circ is the symbol for the multiplication operator, and wherein any non-integer product of x_a and y is rounded down to the nearest integer prior to subtracting it from x_a .

By way of example, a polypeptide sequence of the present invention may be identical to the reference sequence of SEQ ID NO:2, that is it may be 100% identical, or it may include up to a certain integer number of amino acid alterations as compared to the reference sequence such that the percent identity is less than 100% identity. Such alterations are selected from the group consisting of at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion, and wherein said alterations may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between those terminal positions, interspersed either individually among the amino acids in the reference sequence or in one or more contiguous groups within the reference sequence. The number of amino acid alterations for a given % identity is determined by multiplying the total number of amino acids in SEQ ID NO:2 by the integer defining the percent identity divided by 100 and then subtracting that product from said total number of amino acids in SEQ ID NO:2, or:

$$n_a \leq x_a - (x_a \circ y),$$

wherein n_a is the number of amino acid alterations, x_a is the total number of amino acids in SEQ ID NO:2, y is, for instance 0.70 for 70%, 0.80 for 80%, 0.85 for 85% etc., and \circ is the symbol for the multiplication operator, and wherein any non-integer product of x_a and y is rounded down to the nearest integer prior to subtracting it from x_a .

"Individual(s)," when used herein with reference to an organism, means a multicellular eukaryote, including, but not limited to a metazoan, a mammal, an ovid, a bovid, a simian, a primate, and a human.

"Isolated" means altered "by the hand of man" from its natural state, *i.e.*, if it occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living organism is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein. Moreover, a polynucleotide or polypeptide that is introduced into an organism by transformation, genetic manipulation or by any other recombinant method is "isolated" even if it is still present in said organism, which organism may be living or non-living.

"Polynucleotide(s)" generally refers to any polyribonucleotide or polydeoxyribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA including single and double-stranded regions.

"Variant" refers to a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino

acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques or by direct synthesis.

"Disease(s)" means any disease caused by or related to infection by a bacteria, including, for example, otitis media in infants and children, pneumonia in elderlies, sinusitis, nosocomial infections and invasive diseases, chronic otitis media with hearing loss, fluid accumulation in the middle ear, auditive nerve damage, delayed speech learning, infection of the upper respiratory tract and inflammation of the middle ear.

EXAMPLES:

The examples below are carried out using standard techniques, which are well known and routine to those of skill in the art, except where otherwise described in detail. The examples are illustrative, but do not limit the invention.

Example 1: DNA sequencing of the BASB112 gene from *Moraxella catarrhalis* strain ATCC 43617.

A: BASB112 in *Moraxella catarrhalis* strain.

The BASB112 gene of SEQ ID NO:1 is from *Moraxella catarrhalis* strain ATCC 43617. The translation of the BASB112 polynucleotide sequence is showed in SEQ ID NO:2.

B: BASB112 in *Moraxella Catarrhalis* strain 43617.

The sequence of the BASB112 gene was confirmed in *Moraxella Catarrhalis* strain ATCC 43617. For this purpose, plasmid DNA (see example 2A) containing the gene region encoding the mature BASB112 from *Moraxella Catarrhalis*. strain ATCC 43617 used as a PCR template. This material was then submitted to Polymerase Chain Reaction DNA amplification using primers *Moraxella catarrhalis* MCD15b-01 (5'-AGG CAG AGG GAA TTC ATG CAA AAG ATG ACT TAC AAA GC -3') [SEQ ID NO:5] and reverse MCD15b-02 (5'-AGG CAG AGG GTC GAC TTA ATG GTG ATG GTG ATG GTG ATG GCG AGT ATC AGG CAA GCT GTC ACG AC -3') [SEQ ID NO:6] specific for the BASB112 gene. The PCR amplicon was then submitted to DNA sequencing using the Big Dyes kit (Applied biosystems) and analyzed on a ABI 373/A DNA sequencer in the conditions described by the supplier. As a result, the polynucleotide and deduced polypeptide sequences, referred to as SEQ ID NO:3 and SEQ ID NO:4 respectively, were obtained. These sequences do not comprise the signal sequence as the signal sequence was from the plasmid.

Using the MegAlign program from the DNASTAR software package, an alignment of the polynucleotide sequences of SEQ ID NO:1 and 3 was performed, and is displayed in Figure 1; a pairwise comparison of identities shows that the two BASB112 polynucleotide gene sequences are 99.9% identical in the region coding for the mature protein. Using the same MegAlign program, an alignment of the polypeptide sequences of SEQ ID NO:2 and 4 was performed, and is displayed in Figure 2; a pairwise comparison of identities shows that the two BASB112 protein sequences are 99.9% identical in the region of the mature protein.

Example 2: Construction of Plasmid to Express Recombinant BASB112

A: Cloning of BASB112.

The *EcoRI* and *SaII* restriction sites engineered into the MC-Lip3-Fn/t-RI (5'- AGG CAG AGG GAA TTC ATG CAA AAG ATG ACT TAC AAA GC -3') [SEQ ID NO:5] forward and MC-Lip3-RCh/t-Sal (5'- AGG CAG AGG GTC GAC TTA ATG GTG ATG GTG ATG GTG ATG GCG AGT ATC AGG CAA GCT GTC ACG AC - 3') [SEQ ID NO:6] reverse amplification primers, respectively, permitted directional cloning a PCR product into the *E.coli* expression plasmid pTLZ2 such that a mature BASB112 protein could be expressed as a fusion protein containing a (His)₆ affinity chromatography tag at the C-terminus. The BASB112 PCR product was purified from the amplification reaction using silica gel-based spin columns (QiaGen) according to the manufacturers instructions. To produce the required *EcoRI* and *SaII* termini necessary for cloning, purified PCR product was sequentially digested to completion with *EcoRI* and *SaII* restriction enzymes as recommended by the manufacturer (Life Technologies). Following the first restriction digestion, the PCR product was purified via spin column as above to remove salts and eluted in sterile water prior to the second enzyme digestion. The digested DNA fragment was again purified using silica gel-based spin columns prior to ligation with the pTLZ2 plasmid.

B: Production of Expression Vector.

To prepare the expression plasmid pTLZ2 for ligation, it was similarly digested to completion with both *EcoRI* and *SaI* and then treated with calf intestinal phosphatase (CIP, ~0.02 units / pmole of 5' end, Life Technologies) as directed by the manufacturer to prevent self ligation. An approximately 5-fold molar excess of the digested fragment to the prepared vector was used to program the ligation reaction. A standard ~20 µl ligation reaction (~16°C, ~16 hours), using methods well known in the art, was performed using T4 DNA ligase (~2.0 units / reaction, Life Technologies). An aliquot of the ligation (~5 µl) was used to transform electro-competent JM109 cells according to methods well known in the art. Following a ~2-3 hour outgrowth period at 37°C in ~1.0 ml of LB broth, transformed cells were plated on LB agar plates containing ampicillin (100 µg/ml). Antibiotic was included in the selection. Plates were incubated overnight at 37°C for ~16 hours. Individual ampicillin resistant colonies were picked with sterile toothpicks and used to "patch" inoculate fresh LB ApR plates as well as a ~1.0 ml LB ApR broth culture. Both the patch plates and the broth culture were incubated overnight at 37°C in either a standard incubator (plates) or a shaking water bath. A whole cell-based PCR analysis was employed to verify that transformants contained the BASB112 DNA insert. Here, the ~1.0 ml overnight LB Ap broth culture was transferred to a 1.5 ml polypropylene tube and the cells collected by centrifugation in a Beckmann microcentrifuge (~3 min., room temperature, ~12,000 X g). The cell pellet was suspended in ~200µl of sterile water and a ~10 µl aliquot used to program a ~50µl final volume PCR reaction containing both BASB112 forward and reverse amplification primers. Final concentrations of the PCR reaction components were essentially the same as those specified in example 2 except ~5.0 units of *Taq* polymerase was used. The initial 95°C denaturation step was increased to 3 minutes to ensure thermal disruption of the bacterial cells and liberation of plasmid DNA. An ABI Model 9700 thermal cycler and a 32 cycle, three-step thermal amplification profile, i.e. 95°C, 45sec; 55-58°C, 45sec, 72°C, 1min., were used to amplify the BASB112

fragment from the lysed transformant samples. Following thermal amplification, a ~20µl aliquot of the reaction was analyzed by agarose gel electrophoresis (0.8 % agarose in a Tris-acetate-EDTA (TAE) buffer). DNA fragments were visualized by UV illumination after gel electrophoresis and ethidium bromide staining. A DNA molecular size standard (1 Kb ladder, Life Technologies) was electrophoresed in parallel with the test samples and was used to estimate the size of the PCR products. Transformants that produced the expected size PCR product were identified as strains containing a BASB112 expression construct. Expression plasmid containing strains were then analyzed for the inducible expression of recombinant BASB112.

C: Expression Analysis of PCR-Positive Transformants.

For each PCR-positive transformant identified above, ~5.0 ml of LB broth containing ampicillin (Ap) (100 µg/ml) was inoculated with cells from the patch plate and grown overnight at 37 °C with shaking (~250 rpm). An aliquot of the overnight seed culture (~1.0 ml) was inoculated into a 125 ml erlenmeyer flask containing ~25 of LB Ap broth and grown at 37 °C with shaking (~250 rpm) until the culture turbidity reached O.D.600 of ~0.5, i.e. mid-log phase (usually about 1.5 - 2.0 hours). At this time approximately half of the culture (~12.5 ml) was transferred to a second 125 ml flask and expression of recombinant BASB112 protein induced by the addition of IPTG (1.0 M stock prepared in sterile water, Sigma) to a final concentration of 1.0 mM. Incubation of both the IPTG-induced and non-induced cultures continued for an additional ~4 hours at 37 °C with shaking. Samples (~1.0 ml) of both induced and non-induced cultures were removed after the induction period and the cells collected by centrifugation in a microcentrifuge at room temperature for ~3 minutes. Individual cell pellets were suspended in ~50µl of sterile water, then mixed with an equal volume of 2X Laemmli SDS-PAGE sample buffer containing 2-mercaptoethanol, and placed in boiling water bath for ~3 min to denature protein. Equal volumes (~15µl) of both the crude IPTG-induced and the non-induced cell lysates were loaded onto duplicate 12% Tris/glycine polyacrylamide gel (1 mm thick Mini-gels, Novex). The induced and non-induced

lysate samples were electrophoresed together with prestained molecular weight markers (SeeBlue, Novex) under conventional conditions using a standard SDS/Tris/glycine running buffer (BioRad). Following electrophoresis, one gel was stained with commassie brilliant blue R250 (BioRad) and then destained to visualize novel BASB112 IPTG-inducible protein(s). The second gel was electroblotted onto a PVDF membrane (0.45 micron pore size, Novex) for ~2 hrs at 4 °C using a BioRad Mini-Protean II blotting apparatus and Towbin's methanol (20 %) transfer buffer. Blocking of the membrane and antibody incubations were performed according to methods well known in the art. A monoclonal anti-RGS (His)₃ antibody, followed by a second rabbit anti-mouse antibody conjugated to HRP (QiaGen), was used to confirm the expression and identity of the BASB112 recombinant protein. Visualization of the anti-His antibody reactive pattern was achieved using either an ABT insoluble substrate or using Hyperfilm with the Amersham ECL chemiluminescence system.

Example 3: Production of Recombinant BASB112

Bacterial strain

A recombinant expression strain of *E. coli* JM109 containing a plasmid (pTLZ2) encoding BASB112 from *M. catarrhalis*. was used to produce cell mass for purification of recombinant protein. The expression strain was cultivated on LB agar plates containing 100µg/ml Ap to ensure that the pTLZ2 was maintained. For cryopreservation at -80 °C, the strain was propagated in LB broth containing the same concentration of antibiotics then mixed with an equal volume of LB broth containing 30% (w/v) glycerol.

Media

The fermentation medium used for the production of recombinant protein consisted of 2X YT broth (Difco) containing 100µg/ml Ap. Antifoam was added to medium for the fermentor at 0.25 ml/L (Antifoam 204, Sigma). To induce expression of the BASB112 recombinant protein, IPTG (Isopropyl β-D-Thiogalactopyranoside) was added to the

fermentor (1 mM, final).

Fermentation

A 500-ml erlenmeyer seed flask, containing 50ml working volume, was inoculated with 0.3 ml of rapidly thawed frozen culture, or several colonies from a selective agar plate culture, and incubated for approximately 12 hours at $37 \pm 1^\circ\text{C}$ on a shaking platform at 150 rpm (Innova 2100, New Brunswick Scientific). This seed culture was then used to inoculate a 5-L working volume fermentor containing 2X YT broth and both Ap antibiotics. The fermentor (Bioflo 3000, New Brunswick Scientific) was operated at $37 \pm 1^\circ\text{C}$, 0.2 - 0.4 VVM air sparge, 250 rpm in Rushton impellers. The pH was not controlled in either the flask seed culture or the fermentor. During fermentation, the pH ranged 6.5 to 7.3 in the fermentor. IPTG (1.0 M stock, prepared in sterile water) was added to the fermentor when the culture reached mid-log of growth (~ 0.7 O.D.600 units). Cells were induced for 2 - 4 hours then harvested by centrifugation using either a 28RS Heraeus (Sepatech) or RC5C superspeed centrifuge (Sorvall Instruments). Cell paste was stored at -20°C until processed.

Chemicals and Materials

Imidazole and Triton X-100 were purchased from Merck. Guanidine hydrochloride was from Fluka. Aprotinin was obtained from Sigma Chemical Company. Urea and AEBSF were from ICN-Biochemicals. All other chemicals were reagent grade or better.

Ni-NTA Superflow resin and penta-His Antibody, BSA free were obtained from QiaGen. MicroBCA assay was obtained from Pierce; Amicon 3 filters from Millipore. Dialysis membrane (MWCO12-14000) were from MFPI, USA. Molecular mass marker (BenchMark ladder) was from Life-technologies.

Example 4: Purification of recombinant BASB112 from *E. coli*

Extraction–Purification

Cell paste from 750 ml IPTG induced culture (~4 hours, OD₆₂₀= 0.5) was resuspended in 60 ml of phosphate buffer pH 7.5 containing 1mM AEBSF and 1mM Aprotinin as protease inhibitors. Cells were lysed in a cell disruptor. Lysate was centrifuged at 27,000g for 20 minutes. Pellet was washed once with phosphate buffer pH 7.5 and centrifuged again at 27,000g for 20 minutes. Pellet was suspended in 100 mM NaH₂PO₄, 10 mM Tris-HCl buffer pH 8 containing 6M Guanidium Chloride (buffer A) and left for 1 hour at room temperature. Total extract was centrifuged at 27,000g for 20 minutes. Supernatant was incubated for 1 hour at room temperature with Ni-NTA superflow resin equilibrated in buffer A. Resin was washed twice with 100 mM NaH₂PO₄, 10 mM Tris-HCl buffer pH 6.3, containing 8M Urea (buffer B). Elution was performed successively with buffer B adjusted to pH 5.9 then pH 4.5. Fractions containing BASB112 antigen were neutralized with 25% volume of 0.2M phosphate buffer pH 7.5. Pooled fractions were dialysed successively against 100 mM NaH₂PO₄ containing 8M Urea, then 4M Urea, then 2M Urea and finally against PBS pH 7.4 containing 0.1% Triton X-100.

Purified BASB112 antigen was quantified using Micro BCA assay reagent. 1.7 mg of purified antigen were obtained , at a final concentration of 220 µg/ml.

As shown in figure 3-A, purified BASB112 protein appeared in SDS-PAGE analysis as a major band migrating at around 15 kDa (estimated relative molecular mass). Purity was estimated to more than 90 %. BASB112 protein was reactive against a mouse monoclonal antibody raised against the 6-Histidine motif (figure 3-B).

Example 5: Production of Antisera to Recombinant BASB112

Polyvalent antisera directed against the BASB112 protein are generated by vaccinating rabbits with the purified recombinant BASB112 protein. Animals are bled prior to the first immunization ("pre-bleed") and after the last immunization.

Anti-BASB112 protein titres are measured by an ELISA using purified recombinant BASB112 protein. The titre is defined as mid-point titers calculated by 4-parameter logistic model using the XL Fit software.

The antisera are also used as the first antibody to identify the protein in a western blot as described in example 7 below. The western-blot is used to demonstrate the presence of anti-BASB112 antibody in the sera of immunized animals.

Example 6: Immunological characterization: Surface exposure of BASB112

Anti-BASB112 protein titres are determined by an ELISA using formalin-killed whole cells of *Moraxella catarrhalis*. The titre is defined as mid-point titers calculated by 4-parameter logistic model using the XL Fit software.

The titre observed with the rabbit or mouse immune sera demonstrates that the BASB112 protein is present at the surface of *M. catarrhalis* cells.

Example 7. Immunological Characterisation: Western Blot Analysis

Several strains of *M. catarrhalis* including ATCC 43617, as well as clinical isolates from various geographic regions, are grown on Muller Hinton agar plates for 24 hours at 36°C. Several colonies are used to inoculate broth. Cultures are grown until the A620 is approximately 0.6 and cells are collected by centrifugation. Cells are then concentrated and solubilized in PAGE sample buffer. The solubilized cells are then resolved on 4-20% polyacrylamide gels and the separated proteins are electrophoretically transferred to PVDF membranes. The PVDF membranes are then pretreated with saturation buffer. All subsequent incubations are carried out using this pretreatment buffer.

PVDF membranes are incubated with preimmune serum or rabbit or mouse immune

seru. PVDF membranes are then washed.

PVDF membranes are incubated with biotin-labeled sheep anti-rabbit or mouse Ig. PVDF membranes are then washed 3 times with wash buffer, and incubated with streptavidin-peroxydase. PVDF membranes are then washed 3 times with wash buffer and developed with 4-chloro-1-naphtol.

Detection of a protein corresponding to BASB112 expected molecular weight that is reactive with the antisera is used to demonstrate that this protein is produced by and conserved in all *Moraxella* strains tested.

Example 8: Immunological characterization: Bactericidal Activity

Complement-mediated cytotoxic activity of anti-BASB112 antibodies is examined to determine the vaccine potential of BASB112 protein antiserum that is prepared as described above. The activities of the pre-immune serum and the anti-BASB112 antiserum in mediating complement killing of *M. catarrhalis* are examined.

Strains of *M. catarrhalis* are grown on plates. Several colonies are added to liquid medium. Cultures are grown and collected until the A620 is approximately 0.4. After one wash step, the pellet is suspended and diluted.

Preimmune sera and the anti-BASB112 sera is deposited into the first well of a 96-wells plate and serial dilutions are deposited in the other wells of the same line. Live diluted *M. catarrhalis* is subsequently added and the mixture is incubated. Complement is added into each well at a working dilution defined beforehand in a toxicity assay.

Each test includes a complement control (wells without serum containing active or inactivated complement source), a positive control (wells containing serum with a know titer of bactericidal antibodies), a culture control (wells without serum and complement) and a serum control (wells without complement).

Bactericidal activity of rabbit or mice antiserum (50% killing of homologous strain) is

WO 01/00835

PCT/EP00/05849

measured.

Example 9: Presence of Antibody to BASB112 in Human Convalescent Sera

Western blot analysis of purified recombinant BASB112 is performed as described in Example 7 above, except that a pool of human sera from children infected by *M. catarrhalis* is used as the first antibody preparation. This is used to show that antisera from naturally infected individuals react to the purified recombinant protein.

Example 10: Efficacy of BASB112 vaccine: enhancement of lung clearance of *M. catarrhalis* in mice.

This mouse model is based on the analysis of the lung invasion by *M. catarrhalis* following a standard intranasal challenge to vaccinated mice.

Groups of mice are immunized with BASB112 vaccine. After the booster, the mice are challenged by instillation of bacterial suspension into the nostril under anaesthesia. Mice are killed between 30 minutes and 24 hours after challenge and the lungs are removed aseptically and homogenized individually. The log₁₀ weighted mean number of CFU/lung is determined by counting the colonies grown on agar plates after plating of dilutions of the homogenate. The arithmetic mean of the log₁₀ weighted mean number of CFU/lung and the standard deviations are calculated for each group. Results are analysed statistically.

In this experiment groups of mice are immunized either with BASB112 or with a killed whole cells (kwc) preparation of *M. catarrhalis* or sham immunized.

This is used to show that the kwc preparation and the BASB112 vaccine induce significant lung clearance as compared to the control group.

Example 11: Inhibition of *M. catarrhalis* adhesion onto cells by anti-BASB112 antiserum.

This assay measures the capacity of anti BASB112 sera to inhibit the adhesion of

Moraxella bacteria to epithelial cells. This activity could prevent colonization of f.i. the nasopharynx by *Moraxella*.

One volume of bacteria is incubated on ice with one volume of pre-immune or anti-BASB112 immune serum dilution. This mixture is subsequently added in the wells of a 24 well plate containing a confluent cells culture that is washed once with culture medium to remove traces of antibiotic. The plate is centrifuged and incubated.

Each well is then gently washed. After the last wash, sodium glycocholate is added to the wells. After incubation, the cell layer is scraped and homogenised. Dilutions of the homogenate are plated on agar plates and incubated. The number of colonies on each plate is counted and the number of bacteria present in each well calculated.

This is used to show that bacteria incubated with anti-BASB112 antiserum are inhibited in their adherence capacity to the Hep-2 cells.

Deposited materials

A deposit containing a *Moraxella catarrhalis* Catlin strain has been deposited with the American Type Culture Collection (herein "ATCC") on June 21, 1997 and assigned deposit number 43617. The deposit was described as *Branhamella catarrhalis* (Frosch and Kolle) and is a freeze-dried, 1.5-2.9 kb insert library constructed from *M. catarrhalis* isolate obtained from a transtracheal aspirate of a coal miner with chronic bronchitis. The deposit is described in Antimicrob. Agents Chemother. 21: 506-508 (1982).

The *Moraxella catarrhalis* strain deposit is referred to herein as "the deposited strain" or as "the DNA of the deposited strain."

The deposited strain contains a full length BASB112 gene.

A deposit of the vector pMC-D15 consisting of *Moraxella catarrhalis* DNA inserted in pQE30 has been deposited with the American Type Culture Collection (ATCC) on February 12 1999 and assigned deposit number 207105.

The sequence of the polynucleotides contained in the deposited strain / clone, as well as the amino acid sequence of any polypeptide encoded thereby, are controlling in the event of any conflict with any description of sequences herein.

The deposit of the deposited strains have been made under the terms of the Budapest Treaty on the International Recognition of the Deposit of Micro-organisms for Purposes of Patent Procedure. The deposited strains will be irrevocably and without restriction or condition released to the public upon the issuance of a patent. The deposited strains are provided merely as convenience to those of skill in the art and are not an admission that a deposit is required for enablement, such as that required under 35 U.S.C. §112.

Applicant's or agent's
file reference

FB/BM45397

International application no.

PCT/EP 00 / 05849

INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description
on page 65 . line 1-29

B. IDENTIFICATION OF DEPOSIT

Further deposits are identified on an additional sheet ☐

Name of depositary institution

AMERICAN TYPE CULTURE COLLECTION

Address of depositary institution (including postal code and country)

10801 UNIVERSITY BLVD, MANASSAS, VIRGINIA 20110-2209,
UNITED STATES OF AMERICADate of deposit 21 June 1997 (21.06.97)
and 12 February 1999 (12.02.99)Accession Number
43617 and 207105

C. ADDITIONAL INDICATIONS (leave blank if not applicable)

This information is continued on an additional sheet ☐

In respect of those designations where a European Patent is sought, a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European Patent or until the date on which the application has been refused or withdrawn, only by issue of such a sample to an expert nominated by the person requesting the sample.

D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)

E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)

The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")

For receiving Office use only

☒ This sheet was received with the international application

23 JUN 2000

Authorized officer

Y. Marinus-v.d. Nouweland

For International Bureau use only

☐ This sheet was received by the International Bureau on:

Authorized officer

SEQUENCE INFORMATION

BASB112 Polynucleotide and Polypeptide Sequences

SEQ ID NO:1

Moraxella catarrhalis BASB112 polynucleotide sequence from strain ATCC43617

ATGCAAAAGATGACTTACAAAGCTATCGGCTTAGGCTTGGTGCTTGCTTTGACTGGCTGCAGTGTCTTTCGGGTGTATAC
CATCGACTTACCTCAAGGTACGCCACTATCTGCCGCCCAAGCCAATCAAATACAAGTGGGCATGAATGCCAACCAAGTGC
TATATTTACTTGGCTCGCCTGCCTTGCGTGACACCCTAGCACCAAACCGCTGGGATTATATTTATGACCATACCGCACGC
ACGGATGGCAAACGCCAAAAAAATCTGACATTAAAAACGCATCGCAATACTTGAGCATTCAATTTAATAACCAAGGCAT
CGTGACACGCATTGACGGTCGTGACAGCTTGCTGATACTCGCCATTAA

SEQ ID NO:2

Moraxella catarrhalis BASB112 polypeptide sequence deduced from the polynucleotide of
SeQ ID NO:1

MQKMTYKAIGLGLVLALTGCSVFRVYTIDLPQGTPLSAAQANQIQVGMNANQVLYLLGSPALRDTLAPNRWDYIYDHTAR
TDGKRQKSDIKNASQYLSIHFNQGIVTRIDGRDSLDPDTRH

SEQ ID NO:3

Moraxella catarrhalis BASB112 polynucleotide sequence from strain ATCC43617

ATGCAAAAGATGACTTACAAAGCTATcgGCTTAGGCTTGGTGCTTGCTTTGACTGGCTGCAGTGTCTTTCGGGTGTATAC
CATCGACTTACCTcAAGGTACGCCACTATCTGCCGCCCAAGCCAATCAAATACAAGTGGGCATGAATGCCAACCAAGTGC
TATATTTACTTGGCTCGCCTGCCTTGCGTGACACCCTAGCACCAAACCGCTGGGATTATATTTATGACCATACCGCAGGC
ACGGATGGCAAACGCCGAAAAAAATCTGACATTAAAAACGCATCGCAATACTTGAGCATTCAATTTAATAACCAAGGCAT
CGTGACACGCATTGACGGTCGTGACAGCTTGCTGATACTCGCCAT

SEQ ID NO:4

Moraxella catarrhalis BASB112 polypeptide sequence deduced from the polynucleotide of
SeQ ID NO:3

MQKMTYKAIGLGLVLALTGCSVFRVYTIDLPQGTPLSAAQANQIQVGMNANQVLYLLGSPALRDTLAPNRWDYIYDHTAG
TDGKRKSDIKNASQYLSIHFNQGIVTRIDGRDSLDPDTRH

SEQ ID NO:5

AGG CAG AGG GAA TTC ATG CAA AAG ATG ACT TAC AAA GC

SEQ ID NO:6

AGG CAG AGG GTC GAC TTA ATG GTG ATG GTG ATG GTG ATG GCG
AGT ATC AGG CAA GCT GTC ACG AC

CLAIMS:

1. An isolated polypeptide comprising an amino acid sequence which has at least 85% identity to the amino acid sequence selected from the group consisting of: SEQ ID NO:2 and SEQ ID NO:4, over the entire length of SEQ ID NO:2 or SEQ ID NO:4 respectively.
2. An isolated polypeptide as claimed in claim 1 in which the amino acid sequence has at least 95% identity to the amino acid sequence selected from the group consisting of: SEQ ID NO:2 and SEQ ID NO:4, over the entire length of SEQ ID NO:2 or SEQ ID NO:4 respectively.
3. The polypeptide as claimed in claim 1 comprising the amino acid sequence selected from the group consisting of: SEQ ID NO:2 and SEQ ID NO:4.
4. An isolated polypeptide of SEQ ID NO:2 or SEQ ID NO:4.
5. An immunogenic fragment of the polypeptide as claimed in any one of claims 1 to 4 in which the immunogenic activity of said immunogenic fragment is substantially the same as the polypeptide of SEQ ID NO:2 or SEQ ID NO:4.
6. A polypeptide as claimed in any of claims 1 to 5 wherein said polypeptide is part of a larger fusion protein.
7. An isolated polynucleotide encoding a polypeptide as claimed in any of claims 1 to 6.
8. An isolated polynucleotide comprising a nucleotide sequence encoding a polypeptide that has at least 85% identity to the amino acid sequence of SEQ ID NO:2 or 4 over the entire length of SEQ ID NO:2 or 4 respectively; or a nucleotide sequence complementary to said isolated polynucleotide.

9. An isolated polynucleotide comprising a nucleotide sequence that has at least 85% identity to a nucleotide sequence encoding a polypeptide of SEQ ID NO:2 or 4 over the entire coding region; or a nucleotide sequence complementary to said isolated polynucleotide.
10. An isolated polynucleotide which comprises a nucleotide sequence which has at least 85% identity to that of SEQ ID NO:1 or 3 over the entire length of SEQ ID NO:1 or 3 respectively; or a nucleotide sequence complementary to said isolated polynucleotide.
11. The isolated polynucleotide as claimed in any one of claims 7 to 10 in which the identity is at least 95% to SEQ ID NO:1 or 3.
12. An isolated polynucleotide comprising a nucleotide sequence encoding the polypeptide of SEQ ID NO:2 or SEQ ID NO:4.
13. An isolated polynucleotide comprising the polynucleotide of SEQ ID NO:1 or SEQ ID NO:3.
14. An isolated polynucleotide comprising a nucleotide sequence encoding the polypeptide of SEQ ID NO:2, SEQ ID NO:4 obtainable by screening an appropriate library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO:1 or SEQ ID NO:3 or a fragment thereof.
15. An expression vector or a recombinant live microorganism comprising an isolated polynucleotide according to any one of claims 7 - 14.
16. A host cell comprising the expression vector of claim 15 or a subcellular fraction or a membrane of said host cell expressing an isolated polypeptide comprising an amino acid

sequence that has at least 85% identity to the amino acid sequence selected from the group consisting of: SEQ ID NO:2 and SEQ ID NO:4.

17. A process for producing a polypeptide of claims 1 to 6 comprising culturing a host cell of claim 16 under conditions sufficient for the production of said polypeptide and recovering the polypeptide from the culture medium.
18. A process for expressing a polynucleotide of any one of claims 7 – 14 comprising transforming a host cell with the expression vector comprising at least one of said polynucleotides and culturing said host cell under conditions sufficient for expression of any one of said polynucleotides.
19. A vaccine composition comprising an effective amount of the polypeptide of any one of claims 1 to 6 and a pharmaceutically acceptable carrier.
20. A vaccine composition comprising an effective amount of the polynucleotide of any one of claims 7 to 14 and a pharmaceutically effective carrier.
21. The vaccine composition according to either one of claims 19 or 20 wherein said composition comprises at least one other *Moraxella catarrhalis* antigen.
22. An antibody immunospecific for the polypeptide or immunological fragment as claimed in any one of claims 1 to 6.
23. A method of diagnosing a *Moraxella catarrhalis* infection, comprising identifying a polypeptide as claimed in any one of claims 1 - 6, or an antibody that is immunospecific for said polypeptide, present within a biological sample from an animal suspected of having such an infection.

24. Use of a composition comprising an immunologically effective amount of a polypeptide as claimed in any one of claims 1 – 6 in the preparation of a medicament for use in generating an immune response in an animal.
25. Use of a composition comprising an immunologically effective amount of a polynucleotide as claimed in any one of claims 7 - 14 in the preparation of a medicament for use in generating an immune response in an animal.
26. A therapeutic composition useful in treating humans with *Moraxella catarrhalis* disease comprising at least one antibody directed against the polypeptide of claims 1 – 6 and a suitable pharmaceutical carrier.

Figure 1 : Alignment of the BASB112 polynucleotide sequences.**Identity to SeqID No:1 is indicated by a dot. Gap is indicated by a dash.**

	*	20	*	
Seqid1	:	ATGCAAAAGATGACTTACAAAGCTATCGGC	:	30
Seqid3	:	:	30
	40	*	60	
Seqid1	:	TTAGGCTTGGTGCTTGCTTTGACTGGCTGC	:	60
Seqid3	:	:	60
	*	80	*	
Seqid1	:	AGTGTCTTTCGGGTGTATACCATCGACTTA	:	90
Seqid3	:	:	90
	100	*	120	
Seqid1	:	CCTCAAGGTACGCCACTATCTGCCGCCCAA	:	120
Seqid3	:	:	120
	*	140	*	
Seqid1	:	GCCAATCAAATACAAGTGGGCATGAATGCC	:	150
Seqid3	:	:	150
	160	*	180	
Seqid1	:	AACCAAGTGCTATATTTACTTGGCTCGCCT	:	180
Seqid3	:	:	180

313 Rec'd PCT/PTO 13 DEC 2001

2/4

* 200 *

Seqid1 : GCCTTGCGTGACACCCTAGCACCAAACCGC : 210
Seqid3 : : 210

220 * 240

Seqid1 : TGGGATTATATTTATGACCATACCGCACGC : 240
Seqid3 :G.. : 240

* 260 *

Seqid1 : ACGGATGGCAAACGCCAAAAAAATCTGAC : 270
Seqid3 :G..... : 270

280 * 300

Seqid1 : ATTAAAAACGCATCGCAATACTTGAGCATT : 300
Seqid3 : : 300

* 320 *

Seqid1 : CATTTTAATAACCAAGGCATCGTGACACGC : 330
Seqid3 : : 330

340 * 360

Seqid1 : ATTGACGGTCGTGACAGCTTGCCCTGATACT : 360
Seqid3 : : 360

Seqid1 : CGCCATTAA : 369
Seqid3 : : 366

JC13 Rec'd PCT/PTO 13 DEC 2001

Figure 2 : Alignment of the BASB112 polypeptide sequences.
Identity to SeqID No:2 is indicated by a dot. Gap is indicated by a dash.

```

                *           20           *
Seqid2 : MQKMTYKAIGLGLVLALTGCSVFRVYTIDL : 30
Seqid4 : ..... : 30

                40           *           60
Seqid2 : PQGTPLSAAQANQIQVGMNANQVLYLLGSP : 60
Seqid4 : ..... : 60

                *           80           *
Seqid2 : ALRDTLAPNRWDYIYDHTARTDGKRQKKSD : 90
Seqid4 : .....G.....R.... : 90

                100           *           120
Seqid2 : IKNASQYLSIHFNNQGIVTRIDGRDSL PDT : 120
Seqid4 : ..... : 120


Seqid2 : RH : 122
Seqid4 : .. : 122
```

JC13 Rec'd PCT/PTO 13 DEC 2001

Figure 3-A : Coomassie stained SDS-polyacrylamide gel of purified BASB112

Figure 3-B: Western-blotting of purified BASB112 (anti-His antibody).

Figure 3-A Figure 3-B



C13 Rec'd PCT/PTO 13 DEC 2001

SEQUENCE LISTING

<110> SmithKline Beecham Biologicals S.A.

<120> Novel compounds

<130> BM45397

<160> 6

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 369

<212> DNA

<213> *Moraxella catarrhalis*

<400> 1

atgcaaaaga	tgacttacaa	agctatcggc	ttaggcttgg	tgcttgcttt	gactggctgc	60
agtgtctttc	gggtgtatac	catcgactta	cctcaaggta	cgccactatc	tgccgcccaa	120
gccaatcaaa	tacaagtggg	catgaatgcc	aaccaagtgc	tatatttact	tggtcgcct	180
gccttgcgtg	acaccctagc	accaaaccgc	tgggattata	tttatgacca	taccgcacgc	240
acggatggca	aacgcaaaaa	aaaatctgac	attaaaaacg	catcgcaata	cttgagcatt	300
cattttaata	accaaggcat	cgtgacacgc	attgacggtc	gtgacagctt	gcttgatact	360
cgccattaa						369

<210> 2

<211> 122

<212> PRT

<213> *Moraxella catarrhalis*

<400> 2

Met	Gln	Lys	Met	Thr	Tyr	Lys	Ala	Ile	Gly	Leu	Gly	Leu	Val	Leu	Ala
1			5					10					15		
Leu	Thr	Gly	Cys	Ser	Val	Phe	Arg	Val	Tyr	Thr	Ile	Asp	Leu	Pro	Gln
		20					25					30			
Gly	Thr	Pro	Leu	Ser	Ala	Ala	Gln	Ala	Asn	Gln	Ile	Gln	Val	Gly	Met
		35				40					45				
Asn	Ala	Asn	Gln	Val	Leu	Tyr	Leu	Leu	Gly	Ser	Pro	Ala	Leu	Arg	Asp
	50				55					60					
Thr	Leu	Ala	Pro	Asn	Arg	Trp	Asp	Tyr	Ile	Tyr	Asp	His	Thr	Ala	Arg
65				70					75				80		
Thr	Asp	Gly	Lys	Arg	Gln	Lys	Lys	Ser	Asp	Ile	Lys	Asn	Ala	Ser	Gln
			85				90					95			
Tyr	Leu	Ser	Ile	His	Phe	Asn	Asn	Gln	Gly	Ile	Val	Thr	Arg	Ile	Asp
		100					105					110			
Gly	Arg	Asp	Ser	Leu	Pro	Asp	Thr	Arg	His						
	115						120								

<210> 3

<211> 366

<212> DNA

<213> *Moraxella catarrhalis*

<400> 3

atgcaaaaga	tgacttacaa	agctatcggc	ttaggcttgg	tgcttgcttt	gactggctgc	60
agtgtctttc	gggtgtatac	catcgactta	cctcaaggta	cgccactatc	tgccgcccaa	120


```

gccaatcaaa tacaagtggg catgaatgcc aaccaagtgc tatatttact tggctcgctt 180
gccttgcggtg acaccctagc accaaaccgc tgggattata tttatgacca taccgcaggc 240
acggatggca aacgccgaaa aaaatctgac attaaaaacg catcgcaata cttgagcatt 300
cattttaata accaaggcat cgtgacacgc attgacggtc gtgacagctt gcctgatact 360
cgccat 366

```

<210> 4
 <211> 122
 <212> PRT
 <213> *Moraxella catarrhalis*

```

<400> 4
Met Gln Lys Met Thr Tyr Lys Ala Ile Gly Leu Gly Leu Val Leu Ala
1      5      10      15
Leu Thr Gly Cys Ser Val Phe Arg Val Tyr Thr Ile Asp Leu Pro Gln
20     25     30
Gly Thr Pro Leu Ser Ala Ala Gln Ala Asn Gln Ile Gln Val Gly Met
35     40     45
Asn Ala Asn Gln Val Leu Tyr Leu Leu Gly Ser Pro Ala Leu Arg Asp
50     55     60
Thr Leu Ala Pro Asn Arg Trp Asp Tyr Ile Tyr Asp His Thr Ala Gly
65     70     75     80
Thr Asp Gly Lys Arg Arg Lys Lys Ser Asp Ile Lys Asn Ala Ser Gln
85     90     95
Tyr Leu Ser Ile His Phe Asn Asn Gln Gly Ile Val Thr Arg Ile Asp
100    105    110
Gly Arg Asp Ser Leu Pro Asp Thr Arg His
115    120

```

<210> 5
 <211> 38
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Primer

```

<400> 5
aggcagaggg aattcatgca aaagatgact tacaaagc 38

```

<210> 6
 <211> 65
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Primer

```

<400> 6
aggcagaggg tcgacttaat ggtgatgggtg atggtgatgg cgagtatcag gcaagctgtc 60
acgac 65

```


INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/05849

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/31 C12N15/62 C07K14/21 C07K16/12 A61K39/02
A61K48/00 G01N33/50

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K A61K G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EMBL, STRAND, WPI Data, EPO-Internal, PAJ, BIOSIS, MEDLINE, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	OCHSNER U A ET AL.: "Pseudomonas aeruginosa fur overlaps with a gene encoding a novel outer membrane lipoprotein, OmlA" JOURNAL OF BACTERIOLOGY, vol. 181, no. 4, February 1999 (1999-02), pages 1099-1109, XP002154428 Note: 31 % aa seq identity with SEQ ID NO:2 in 116 aa overlap. abstract	1-5,7-13
A	WO 94 10316 A (UNIV SASKATCHEWAN) 11 May 1994 (1994-05-11) claims 1-11; examples 1-6 --- -/--	1-26

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

1 December 2000

Date of mailing of the international search report

15/12/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

van de Kamp, M

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 00/05849

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
------------	------------------------------------------------------------------------------------	-----------------------

A	<p>MURPHY T F: "Branhamella catarrhalis: epidemiology, surface antigenic structure, and immune response" MICROBIOLOGICAL REVIEWS, vol. 60, no. 2, July 1996 (1996-07), pages 267-279, XP000857203 cited in the application page 271, right-hand column, line 57 -page 273, right-hand column, line 27; figure 2; table 3</p>	1-26
A	<p>WO 98 55606 A (LOOSMORE SHEENA M ;CONNAUGHT LAB (CA); WANG QUIJUN (CA); YANG YAN) 10 December 1998 (1998-12-10) page 3, line 20 -page 11, line 5 example 4</p>	1-26
A	<p>PANDHER K ET AL.: "Genetic and immunologic analyses of PlpE, a lipoprotein important in complement-mediated killing of Pasteurella haemolytica serotype 1" INFECTION AND IMMUNITY, vol. 66, no. 12, December 1998 (1998-12), pages 5613-5619, XP002154430 abstract</p>	19-22
A	<p>GRAM T ET AL.: "Improved diagnostic PCR assay for Actinobacillus pleuropneumoniae based on the nucleotide sequence of an outer membrane lipoprotein" JOURNAL OF CLINICAL MICROBIOLOGY, vol. 36, no. 2, February 1998 (1998-02), pages 443-448, XP002154429 abstract</p>	23

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 00/05849

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9410316 A	11-05-1994	US 5441736 A	15-08-1995
		CA 2148369 A	11-05-1994
		EP 0669984 A	06-09-1995
		US 5891677 A	06-04-1999
WO 9855606 A	10-12-1998	US 5977337 A	02-11-1999
		AU 7753498 A	21-12-1998
		EP 1000144 A	17-05-2000

